



ISSN No: 0975-7384
CODEN(USA): JCPRC5

J. Chem. Pharm. Res., 2011, 3(2):172-181

Study of Molecular Orbitals of Ruthenium (II) Chloride Based on Molecular Mechanics

Gayasuddin Khan^{1*}, R. N. Verma¹ and B. P. Pandey²

¹Department of Physics, K. S. Saket Post Graduate College, Ayodhya, Faizabad, U.P., INDIA

²Department of Physics, Maharani Lal Kunwari Post Graduate College, Balrampur, U.P., INDIA

ABSTRACT

We have studied the molecular orbitals of ruthenium (II) chloride, in order to study the extent of contribution of 4d, 5s and 5p orbitals in the formation of molecular orbitals. The 3D modeling and geometry optimization of the ruthenium (II) chloride have been done by CAChe software using molecular mechanics with EHT option. Eigenvector analysis shows that $4dx^2-y^2$ and $4dxy$ orbitals of ruthenium play a major role in bonding between ruthenium and chloride, 5s orbital is next and 4p orbitals have a negligible role. There is a difference in energy levels of s and p orbitals of chloride are 0.1691 eV. The overlap population analysis shows that the nonbonding orbitals are present in 6th and 7th molecular orbitals. No molecular orbital is formed by only two atomic orbitals. All molecular orbitals have contribution from many atomic orbitals; the difference is only in extent of involvement.

Keywords: Ruthenium (II) chloride, sd hybridization, Population analysis, Overlap population analysis, Eigenvector, Eigenvalues.

INTRODUCTION

In the recent years Landis [1- 4] and others [5, 6] have considered only ns and (n-1)d orbitals as valence orbitals of the transition metals. They have ignored the involvement of np orbitals. It has been shown that in hybridization only s and d orbitals are involved. They have also described the hybridization angles and idealized molecular shapes of sd, sd², sd³, sd⁴ and sd⁵ hybridizations [7-9]. The restriction to valence s and d functions of transition metals suggested by Landis [2-4] means that 12 electrons will fill the transition metal valence shell

rather than the 18 electrons that can be accommodated if np orbitals were also part of the valence shell. This is astonishing in the light of 18e rule of transition metal compounds. As support for the hypothesis of 12 electron valence space, Landis presented the result of DFT calculation of transition metal hydride [2-4]. He also gave the results of an NBO analysis of the transition metal-hydrogen bonds, which show dominantly sd^n hybridized bond orbitals and negligible np participation [2] However, there is a serious technical flaw in the analysis. The NBO method requires preselection of those orbitals, which are considered as valence orbitals, and may become occupied in the population analysis. In this paper we present the calculations of eigenvalues, eigenvector, overlap matrix and population analysis of ruthenium (II) chloride, in order to study the extent of contribution of 4d, 5s and 5p orbitals in the formation of molecular orbitals. Such a quantitative study will provide correct information about the involvement of 5p orbital of ruthenium in bonding.

EXPERIMENTAL SECTION

The study materials of this paper are ruthenium (II) chloride. The 3D modeling and geometry optimization of the ruthenium (II) chloride have been done by CAChe software using molecular mechanics with EHT option. Eigenvalues, eigenvectors and overlap matrix values have been obtained with the same software, using the same option. With the help of these values, eigenvector analysis, magnitude of contribution of atomic orbital in MO formation, population analysis and overlap population analysis have been made and discussed. The theory on which various calculations are made is defined elsewhere [10].

RESULTS AND DISCUSSION

Ruthenium (II) chloride is triatomic molecule, having the following optimized geometry [11-12] as obtained from molecular mechanics [13–16] method.




Figure.1 Optimized geometry of ruthenium (II) chloride.

Table 1. Eigenvector values of molecular orbitals of Ruthenium (II) chloride.

Atom	AOs	Eigenvector values or coefficients of Atomic Orbitals																
	(χ)	MO-1	MO-2	MO-3	MO-4	MO-5	MO-6	MO-7	MO-8	MO-9	MO-10	MO-11	MO-12	MO-13	MO-14	MO-15	MO-16	MO-17
Ru-1	5s	-0.1029	0.0000	0.0966	-0.0000	0.0000	0.0000	0.0000	-0.0000	-0.0000	0.0000	-0.4603	-0.0000	-0.0000	-0.0000	-0.0000	-1.1218	0.0000
	5px	-0.0000	-0.0688	-0.0000	0.0000	-0.0000	0.0000	0.0000	-0.1681	-0.0014	0.0000	0.0000	-0.0000	-0.0000	0.0111	0.0000	0.0000	1.4554
	5py	0.0000	-0.0007	0.0000	0.0000	-0.0000	-0.0000	-0.0000	-0.0018	0.1271	-0.0006	-0.0000	0.0000	0.0000	-1.0225	-0.0003	0.0000	0.0158
	5pz	-0.0000	-0.0000	-0.0000	0.0000	0.0000	0.0000	-0.0000	0.0000	0.0006	0.1271	-0.0000	-0.0000	0.0000	0.0003	-1.0226	-0.0000	-0.0000
	4dx ² -y ²	-0.1049	-0.0000	0.4628	-0.0027	0.0161	-0.0000	-0.4999	0.0000	0.0000	-0.0000	0.6669	0.0003	0.0146	-0.0000	-0.0000	-0.4071	0.0000
	4dz ²	0.0606	-0.0000	-0.2673	0.0000	-0.0000	-0.0000	-0.8660	-0.0000	-0.0000	0.0000	-0.3851	-0.0000	-0.0000	0.0000	0.0000	0.2351	-0.0000
	4dxy	-0.0023	0.0000	0.0100	0.1234	-0.7415	-0.0000	-0.0108	-0.0000	-0.0000	0.0000	0.0145	-0.0131	-0.6721	-0.0000	-0.0000	-0.0088	-0.0000
	4dxz	0.0000	-0.0000	-0.0000	0.7416	0.1234	0.0108	-0.0000	-0.0000	-0.0000	-0.0000	0.0000	0.6722	-0.0131	0.0000	-0.0000	-0.0000	0.0000
	4dyz	0.0000	0.0000	0.0000	0.0080	0.0013	-0.9999	0.0000	-0.0000	0.0000	0.0000	-0.0000	0.0073	-0.0001	-0.0000	0.0000	-0.0000	-0.0000
Cl-2	3s	-0.6626	-0.6829	-0.1572	0.0000	-0.0000	-0.0000	-0.0000	0.0675	0.0000	-0.0000	-0.0028	0.0073	-0.0000	0.0000	0.0000	0.3728	-0.5297
	3px	-0.0142	-0.0187	-0.4916	-0.0007	0.0043	-0.0000	0.0000	0.6403	-0.0074	0.0000	0.2812	-0.0001	-0.0064	-0.0028	-0.0000	-0.6084	0.6516
	3py	-0.0002	-0.0002	-0.0053	0.0660	-0.3966	-0.0000	-0.0000	0.0069	0.6796	-0.0032	0.0030	0.0115	0.5894	0.2616	0.0001	-0.0066	0.0071
	3pz	0.0000	-0.0000	-0.0000	0.3967	0.0660	0.0000	-0.0000	0.0000	0.0032	0.6796	0.0000	-0.5894	0.0115	-0.0001	0.2616	0.0000	-0.0000
Cl-3	3s	-0.6626	0.6829	-0.1572	-0.0000	0.0000	0.0000	-0.0000	-0.0675	0.0000	-0.0000	-0.0028	0.0000	0.0000	0.0000	0.0000	0.3728	0.5297
	3px	0.0142	-0.0187	0.4916	0.0007	-0.0043	-0.0000	0.0000	0.6403	-0.0074	0.0000	-0.2812	0.0001	0.0064	-0.0028	-0.0000	0.6084	0.6516
	3py	0.0002	-0.0002	0.0053	-0.0660	0.3966	-0.0000	-0.0000	0.0069	0.6796	-0.0032	-0.0030	-0.0115	-0.5894	0.2616	0.0001	0.0066	0.0071
	3pz	0.0000	0.0000	-0.0000	-0.3967	-0.0660	0.0000	-0.0000	0.0000	0.0032	0.6796	0.0000	0.5894	-0.0115	-0.0001	0.2616	0.0000	0.0000

Table 2. Overlap matrix (Overlap integrals values) of Ruthenium (II) chloride.

AOs	5s (Ru-1)	5px (Ru-1)	5py (Ru-1)	5pz (Ru-1)	4dx ² -y ² (Ru-1)	4dz ² (Ru-1)	4dxy (Ru-1)	4dxz (Ru-1)	4dyz (Ru-1)	3s (Cl-2)	3px (Cl-2)	3py (Cl-2)	3pz (Cl-2)	3s (Cl-3)	3px (Cl-3)	3py (Cl-3)	3pz (Cl-3)	
5s (Ru-1)	1.0000																	
5px (Ru -1)	-0.0000	1.0000																
5py (Ru -1)	-0.0000	-0.0000	1.0000															
5pz (Ru -1)	0.0000	0.0000	0.0000	1.0000														
4dx ² -y ² (Ru -1)	0.0000	0.0000	-0.0000	0.0000	1.0000													
4dz ² (Ru -1)	-0.0000	-0.0000	-0.0000	0.0000	0.0000	1.0000												
4dxy (Ru -1)	-0.0000	0.0000	0.0000	0.0000	-0.0000	0.0000	1.0000											
4dxz (Ru 1)	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	1.0000										
4dyz (Ru -1)	0.0000	0.0000	0.0000	0.0000	0.0000	-0.0000	0.0000	-0.0000	1.0000									
3s (Cl-2)	0.2270	0.3406	0.0037	0.0000	0.0994	-0.0574	0.0022	0.0000	0.0000	1.0000								
3px (Cl-2)	-0.3205	-0.3862	-0.0060	0.0000	-0.1310	0.0757	-0.0038	0.0000	0.0000	-0.0000	1.0000							
3py (Cl-2)	-0.0035	-0.0060	0.1707	0.0000	-0.0034	0.0008	0.0923	0.0000	0.0000	-0.0000	0.0000	1.0000						
3pz (Cl-2)	0.0000	0.0000	0.0000	0.1708	0.0000	0.0000	0.0000	0.0923	0.0010	0.0000	0.0000	0.0000	1.0000					
3s (Cl-3)	0.2270	-0.3406	-0.0037	0.0000	0.0994	-0.0574	0.0022	0.0000	0.0000	0.0004	-0.0027	-0.0000	0.0000	1.0000				
3px (Cl-3)	0.3205	-0.3862	-0.0060	0.0000	0.1310	-0.0757	0.0038	0.0000	0.0000	-0.0027	-0.0108	-0.0000	0.0000	0.0000	1.0000			
3py (Cl-3)	0.0035	-0.0060	0.1707	0.0000	0.0034	-0.0008	-0.0923	0.0000	0.0000	0.0000	-0.0001	0.0011	0.0000	-0.0000	-0.0000	1.0000		
3pz (Cl-3)	0.0000	0.0000	0.0000	0.1708	0.0000	0.0000	0.0000	-0.0923	-0.0010	0.0000	0.0000	0.0000	0.0011	0.0000	0.0000	0.0000	1.0000	

The MOs of ruthenium (II) chloride are formed by linear combination of 9 ruthenium orbitals and 4 orbitals from chloride as detailed below-

$$\begin{array}{rcl} \text{Ru-1} & = & 5s, 5p_x, 5p_y, 5p_z, 4d_{x^2-y^2}, 4d_{z^2}, 4d_{xy}, 4d_{xz}, 4d_{yz} = 9 \\ \text{Cl-2} & = & 3s, 3p_x, 3p_y, 3p_z = 4 \\ \text{Cl-3} & = & 3s, 3p_x, 3p_y, 3p_z = 4 \\ & & \text{Total} = 17 \end{array}$$

In order to examine the contribution of various atomic orbitals in the formation of molecular orbitals the LCAO has been studied. The 17 AOs give LCAO approximations to the 17 MOs of ruthenium (II) chloride. The various AOs are represented by χ and MOs by ϕ . χ_1 to χ_9 are 5s, 5p_x, 5p_y, 5p_z, 4d_{x²-y²}, 4d_{z²}, 4d_{xy}, 4d_{xz}, 4d_{yz}, respectively are atomic orbitals of ruthenium and χ_{10} to χ_{13} are 3s, 3p_x, 3p_y, 3p_z for Cl-2; χ_{14} to χ_{17} are 3s, 3p_x, 3p_y, 3p_z for Cl-3, respectively are atomic orbitals of chloride.

The eigenvalues of 17 MOs (ϕ_1 to ϕ_{17}) of ruthenium (II) chloride are -0.9810, -0.9696, -0.5934, -0.5824, -0.5824, -0.5476, -0.5476, -0.5326, -0.5271, -0.5271, -0.4986, -0.4726, -0.4726, -0.2118, -0.2118, 0.1413 and 0.6063, respectively. The coefficients of χ are the eigenvector and overlap matrix which has been taken from Table-1 and Table- 2.

In order to examine the extent of involvement of 4d, 5s and 5p orbitals in the formation of molecular orbitals the values of coefficient of these orbitals have been added to see the total involvement in all the eleven MOs. The summation values of 4d_{xy}, 4d_{xz}, 4d_{x²-y²}, 5s, 5p_x, 5p_y, and 5p_z are 0.9025, 0.8758, 1.7533, 0.6598, 0.2383, 0.1302 and 0.1277, respectively. The nonbonding orbitals 4d_{z²} and 4d_{yz} are excluded. It is clearly indicated that the maximum involvement is of 4d_{x²-y²} orbital and the minimum of 5p_z orbital. The involvement of p orbital is negligible. The value of coefficient is between 0.2383 to 0.1277 which is very low in comparison to d orbitals (d_{xy}, d_{xz}, d_{x²-y²}) which is in the range 1.7533 to 0.8758. The value for 5s is 0.6598. The extent of involvement of 4d, 5s and 5p orbitals of ruthenium in the formation of MOs in the chloride is well demonstrated by the graph (Fig-2) drawn between the orbitals and the summation values of their coefficients. The graph showing below clearly shows that the involvement of p orbitals is negligible.




Figure.2 Trend of extent of involvement of metal orbital in the formation of MOs of CoCl₂

Population Analysis:

The contribution of electrons in each occupied MO is calculated by using the population analysis method, introduced by Mulliken. This method apportions the electrons of n-electron molecule into net population n_r in the basis function χ_r .

Let there be n_i electrons in the MO ϕ_i ($n_i = 0, 1, 2$) and let $n_{r,i}$ symbolize the contribution of electrons in the MO ϕ_i to the net population in χ_r . We have

$$n_{r,i} = n_i c_{ri}^2 \quad (1)$$

where, c_{ri} is the coefficient of atomic orbitals for the i^{th} MO ($r=1-17$).

Equation- 1 has been solved for 22 electrons of 11 molecular orbitals. Two electrons in the 1st MO to 11th MO have been considered. The six molecular orbitals having no electron are left over. The data relating to c_{ri} have been taken from Table 1. The results of solution of equation-1 are included in Table 3 which enlists the contribution of electrons in molecular orbitals under two sections- major and intermediate. It is evident that major contribution is from 4d and 5s orbital. The p orbitals have negligible contribution. The details of contribution are as below.

Table 3: Contribution of electrons in MO of Ruthenium (II) chloride.

MO. No	n_i	Major contribution		Minor contribution	
		Basis function (χ_r)	$n_{r,i} = n_i c_{ri}^2$	Basis function (χ_r)	$n_{r,i} = n_i c_{ri}^2$
1	2	5s (Ru 1)	0.0211	4dz ²	0.0073
		4dx ² -y ² (Ru 1)	0.0220		
		3s (Cl 2)	0.8780		
2	2	3s (Cl 3)	0.8780		
		3s (Cl 2)	0.9327	5px (Ru 1)	0.0094
3	2	3s (Cl 3)	0.9327		
		4dx ² -y ² (Ru 1)	0.4283	5s (Ru 1)	0.0186
		4dz ² (Ru 1)	0.1428	3s (Cl 2)	0.0494
		3px (Cl 2)	0.4833	3s (Cl 3)	0.0494
4	2	3px (Cl 3)	0.4833		
		4dxz (Ru 1)	1.0999	4dxy (Ru 1)	0.0304
		3pz (Cl 2)	0.3147	4py (Cl 2)	0.0087
5	2	3pz (Cl 3)	0.3147	4py (Cl 3)	0.0087
		4dxy (Ru 1)	1.0996	4dxz (Ru 1)	0.0304
		3py (Cl 2)	0.3145	3pz (Cl 2)	0.0087
6	2	3py (Cl 3)	0.3145	3pz (Cl 3)	0.0087
		4dyz (Ru 1)	1.9996		
		4dx ² -y ² (Ru 1)	0.4998		
7	2	4dz ² (Ru 1)	1.4999		
		5px (Ru 1)	0.0565	3s (Cl 2)	0.0091
		3px (Cl 2)	0.8199	3s (Cl 3)	0.0091
8	2	3px (Cl 3)	0.8199		
		5py (Ru 1)	0.0323		
		3py (Cl 2)	0.9237		
9	2	3py (Cl 3)	0.9237		
		5pz (Ru 1)	0.0323		
		3pz (Cl 2)	0.9237		
10	2	3pz (Cl 3)	0.9237		

11	2	5s (Ru 1)	0.4237		
		4dx ² -y ² (Ru 1)	0.8895		
		4dz ² (Ru 1)	0.2866		
		3px (Cl 2)	0.1581		
		3px (Cl 3)	0.1581		

Besides contribution of electrons the Mulliken's method is also used for evaluating overlap population, in order to distinguish bonding, nonbonding and anti bonding molecular orbitals. This method allocates proportionally the overlap population n_{r-s} for all possible pairs of basis functions. This is shown by the equation-2.

$$n_{r-s,i} = n_i (2c_{ri} c_{si} S_{rs}) \quad (2)$$

Where, c_{ri} = the coefficient of atomic orbitals for one atom

c_{si} = the coefficient of atomic orbitals for other atom .

and S_{rs} = the overlap integral between the two AOs (one of an atom and one of other atom).

It is evident from equation-2 that for overlap population analysis of MOs of a molecule, we need eigenvector values (coefficients), values of overlap matrix (overlap integrals) and number of electrons in each MO. The eigenvector and overlap integral values for chloride of ruthenium have been taken from Table-1 and Table-2 respectively and the number of electrons is taken as two for 1st to 11th MOs and zero for 12th to 17th MO. With these values Table 4 is constructed for overlap-population contributions $n_{r-s,i}$ of one molecular orbital. This table has 7 columns, defined as below. There will be 17 such tables for 17 MO but only 11 tables are constructed, because remaining six which have no electrons are left over. In such a way there will be 11 tables.

Column 1 – number of electron n_i

Column 2, 4 – atomic orbitals of ruthenium and chloride.

Column 3 – coefficients of AOs of one atom (c_{ri})

Column 5 – coefficients of AOs of other atom (c_{si})

Column 6 – overlap integral between two AOs of different atoms (S_{rs})

Column 7 – overlap population contribution $n_{r-s,i}$.

The possible overlaps between the various AOs of ruthenium and chloride in each molecular orbital will be 88, as detailed below–

8 overlaps – 5s AO of Ruthenium with 3s, 3px, 3py, 3pz AOs of Cl-2 and Cl-3.

8 overlaps – 5px AO of Ruthenium with 3s, 3px, 3py, 3pz AOs of Cl-2 and Cl-3.

8 overlaps – 5py AO of Ruthenium with 3s, 3px, 3py, 3pz AOs of Cl-2 and Cl-3.

8 overlaps – 5pz AO of Ruthenium with 3s, 3px, 3py, 3pz AOs of Cl-2 and Cl-3.

8 overlaps – 4dx²-y² AO of Ruthenium with 3s, 3px, 3py, 3pz AOs of Cl-2 and Cl-3.

8 overlaps – 4dz² AO of Ruthenium with 3s, 3px, 3py, 3pz AOs of Cl-2 and Cl-3.

8 overlaps – 4dxy AO of Ruthenium with 3s, 3px, 3py, 3pz AOs of Cl-2 and Cl-3.

8 overlaps – 4dxz AO of Ruthenium with 3s, 3px, 3py, 3pz AOs of Cl-2 and Cl-3.

8 overlaps – 4dyz AO of Ruthenium with 3s, 3px, 3py, 3pz AOs of Cl-2 and Cl-3.

4 overlaps – 3s AO of Cl-2 with 3s, 3px, 3py, 3pz AO of Cl-3.

4 overlaps – 3px AO of Cl-2 with 3s, 3px, 3py, 3pz AO of Cl-3.

4 overlaps – 3py AO of Cl-2 with 3s, 3px, 3py, 3pz AO of Cl-3.

4 overlaps – 3pz AO of Cl-2 with 3s, 3px, 3py, 3pz AO of Cl-3.

Total– 88 overlaps

For the study of overlap population we have to construct eleven tables, Having 88 possible overlaps but while building up the table we have dropped the values of zero eigenvector value (Table 1), hence each table of overlap-population contribution differs in its number of orbitals. For obtaining the values of overlap-population contributions ($n_{r-s,i}$) we have to discuss each table separately, but for brevity we here discuss Table 4 for molecular orbital number 1 of ruthenium chloride, which is below:

Table 4. Overlap populations of 1st MO of Ruthenium (II) chloride.						
n_i	AOs	c_{ri}	AOs	c_{si}	S_{rs}	$n_{r-s,i} = n_i(2c_{ri}.c_{si}.S_{rs})$
2	5s(Ru 1)	-0.1029	3s(Cl 2)	-0.6626	0.2270	0.06190
2	5s(Ru 1)	-0.1029	3px(Cl 2)	-0.0142	-0.3205	-0.00187
2	5s(Ru 1)	-0.1029	3py(Cl 2)	-0.0002	-0.0035	0.00000
2	5s(Ru 1)	-0.1029	3s(Cl 3)	-0.6626	0.2270	0.06190
2	5s(Ru 1)	-0.1029	3px(Cl 3)	0.0142	0.3205	-0.00187
2	5s(Ru 1)	-0.1029	3py(Cl 3)	0.0002	0.0035	0.00000
2	4dx ² -y ² (Ru 1)	-0.1049	3s(Cl 2)	-0.6626	0.0994	0.02763
2	4dx ² -y ² (Ru 1)	-0.1049	3px(Cl 2)	-0.0142	-0.1310	-0.00078
2	4dx ² -y ² (Ru 1)	-0.1049	3py(Cl 2)	-0.0002	-0.0034	0.00000
2	4dx ² -y ² (Ru 1)	-0.1049	3s(Cl 3)	-0.6626	0.0994	0.02763
2	4dx ² -y ² (Ru 1)	-0.1049	3px(Cl 3)	0.0142	0.1310	-0.00078
2	4dx ² -y ² (Ru 1)	-0.1049	3py(Cl 3)	0.0002	0.0034	0.00000
2	4dz ² (Ru 1)	0.0606	3s(Cl 2)	-0.6626	-0.0574	0.00921
2	4dz ² (Ru 1)	0.0606	3px(Cl 2)	-0.0142	0.0757	-0.00026
2	4dz ² (Ru 1)	0.0606	3py(Cl 2)	-0.0002	0.0008	0.00000
2	4dz ² (Ru 1)	0.0606	3s(Cl 3)	-0.6626	-0.0574	0.00921
2	4dz ² (Ru 1)	0.0606	3px(Cl 3)	0.0142	-0.0757	-0.00026
2	4dz ² (Ru 1)	0.0606	3py(Cl 3)	0.0002	-0.0008	0.00000
2	4dxy(Ru 1)	-0.0023	3s(Cl 2)	-0.6626	0.0022	0.00001
2	4dxy(Ru 1)	-0.0023	3px(Cl 2)	-0.0142	-0.0038	0.00000
2	4dxy(Ru 1)	-0.0023	3py(Cl 2)	-0.0002	0.0923	0.00000
2	4dxy(Ru 1)	-0.0023	3s(Cl 3)	-0.6626	0.0022	0.00001
2	4dxy(Ru 1)	-0.0023	3px(Cl 3)	0.0142	0.0038	0.00000
2	4dxy(Ru 1)	-0.0023	3py(Cl 3)	0.0002	-0.0923	0.00000
2	3s(Cl 2)	-0.6626	3s(Cl 3)	-0.6626	0.0004	0.00070
2	3s(Cl 2)	-0.6626	3px(Cl 3)	0.0142	0.0027	-0.00010
2	3px(Cl 2)	-0.0142	3s(Cl 3)	-0.6626	-0.0027	-0.00010
2	3px(Cl 2)	-0.0142	3px(Cl 3)	0.0142	-0.0108	0.00000
2	3px(Cl 2)	-0.0142	3py(Cl 3)	0.0002	-0.0001	0.00000
2	3py(Cl 2)	-0.0002	3px(Cl 3)	0.0142	-0.0001	0.00000
2	3py(Cl 2)	-0.0002	3py(Cl 3)	0.0002	0.0011	0.00000
						$\sum n_{r-s,i} = 0.19218$

This table has 31 possible overlaps, out of which 24 provide coefficient values of ruthenium orbitals and 7 for Cl-2, in column 3 that are c_{ri} . Column-5 is for coefficient value c_{si} , for both the chlorines. Up to 24, both the chlorines are involved and for remaining seven only Cl-3. Column-6, is overlap integral S_{rs} and exhibits the magnitude of overlap between the AOs represented in column-2 and 4. The values are self explanatory for indicating the magnitude.

The overlap population analysis also shows negligible involvement of 5p orbitals of ruthenium. It has earlier been suggested that much smaller radius of the 4d orbital than the 5s orbital makes the involvement of 5s orbital dominant contribution in the bonding [17,18]. This hypothesis is the central theme of a recent text book of transition-metal chemistry by Gerloch and Constable [17]. While the importance of the valence ns and (n-1)d functions for the description for transition metal bond is undisputed, the status of the empty np orbital is controversially discussed.

Our results indicate that involvement of np orbital in transition metal bond is negligible and the main role is played by ns and by (n-1)d orbital. Landis [1-4] has also emphatically denied the involvement of np orbital in hybridization. He has supported sd hybridization and support the Landis concept.

The column-7 of Table 4 enlists the values of overlap population, derived from the equation –2. The sum of the values of overlap-populations decides whether the MO in a covalent molecule is bonding, nonbonding or antibonding. If the sum of this inter atomic overlap population contribution is substantially positive, the MO is bonding; if substantially negative, the MO is antibonding and if zero or near zero, the MO is nonbonding. Table 4 indicates that the sum of overlap- population contribution in first MO of RuCl₂ is 0.19218 which is positive indicating or supporting the bonding nature of MO.

Similarly the sum of overlap population for the 11 MO has been worked out and the results are tabulated in Table 5.

Table 5. Nature of occupied MOs			
MO. No	Sum of overlap population contribution ($\sum_{R-s,l}$)		Nature of MOs
1	0.19218	Positive	Bonding
2	0.12479	Positive	Bonding
3	0.22345	Positive	Bonding
4	0.22263	Positive	Bonding
5	-0.21482	Negative	Antibonding
6	0.00000	Zero	Nonbonding
7	0.00000	Zero	Nonbonding
8	0.28493	Positive	Bonding
9	0.06038	Positive	Bonding
10	0.12005	Positive	Bonding
11	0.07327	Positive	Bonding

The overlap population analysis as presented in Table 5 shows that the nonbonding orbitals are present in 6th and 7th molecular orbitals. The difference in positions of nonbonding molecular orbitals prompted us to examine the eigenvalues of ruthenium ion and to compare them with the eigenvalues of the halides. The eigenvalues of the molecular orbitals of the chloride are described above. The nonbonding orbital is degenerate in all the cases. The eigenvector analysis as presented in Table-1 indicates that these orbitals are 4dyz and 4dz².

From the above discussion it is clear that no molecular orbital is formed by only two atomic orbitals. All molecular orbitals have contribution of many basis functions or atomic orbitals;

as a result every molecular orbital has a definite shape having contribution from many basis functions.

CONCLUSION

1. Eigen vector analysis shows that $4dx^2-y^2$ and $4dxy$ orbitals of ruthenium play a major role in bonding between ruthenium and chloride, 5s orbital is next and 4p orbitals have a negligible role. This supports the Landis observation and concept of sd hybridization.
2. s and p orbitals of chloride are involved in bonding with ruthenium. There is a difference in energy levels of s and p orbitals are 0.1691 eV.
3. The overlap population analysis shows that the nonbonding orbitals are present in 6th and 7th molecular orbitals.
4. No molecular orbital is formed by only two atomic orbitals. All molecular orbitals have contribution from many atomic orbitals; the difference is only in extent of involvement.

Acknowledgement

This paper is dedicated to the father of corresponding author.

REFERENCES

- [1] CR Landis; T Cleveland; TK Firman, *J. Am. Chem. Soc.* **1995**, 117, 1859.
- [2] CR Landis; TK Firman; DM Root; T Cleveland; *J. Am. Chem. Soc.* **1998**, 120, 1842.
- [3] CR Landis; T Cleveland; TK Firman, *J. Am. Chem. Soc.* **1998**, 120, 2641.
- [4] TK Firman, CR Landis, *J. Am. Chem. Soc.* **1998**, 120, 12650.
- [5] LG Vanquickenborne; D Pierloot; D Devoghel, *J. Chem. Educ.* **1994**, 71, 469.
- [6] MP Melrose; ER Scerri, *J. Chem. Educ.* **1996**, 73, 498.
- [7] AJ Shortland; G Wilkinson, *J. Chem. Soc. Dalton Trans.* **1973**, 872.
- [8] V Pfennig; K Seppelt, *Science* **1996**, 271, 626.
- [9] M Kaupp, *J. Am. Chem. Soc.* **1996**, 118, 3018.
- [10] PP Singh; PD Sharma, *Russ.J.Coord.Chem.*, 2008, 34, 210-219.
- [11] GE Scuseria, *Chem. Phys. Lett.*, **1995**, 243, 193.
- [12] J Baker; A Kessi, B Delley, *J. Chem. Phys.*, **1996**, 105, 192.
- [13] K Lipkowitz, *J. Chem. Educ.*, **1995**, 72, 1070.
- [14] AK Rappe; CJ Casewit, *Molecular Mechanics Across Chemistry*, University Science Books, **1997**.
- [15] CR Landis, KB Lipkowitz; DB Boyd, *Reviews in Computational Chemistry*, Vol. 6, VCH, **1995**, Chapter 2.
- [16] M Zimmer, *Chem. Rev.*, **1995**, 95, 2629.
- [17] M Gerloch, EC Constable, *Transition-Metal Chemistry*; VCH: Weinheim, **1994**.
- [18] M Gerloch, *Coord. Chem. Rev.*, **1990**, 99, 199.