



Research Article

ISSN : 0975-7384
CODEN(USA) : JCPRC5

An optimization algorithm for solving a class of multiplicative problems

Hongwei Jiao^{1*}, Kun Li² and Jianping Wang¹

¹School of Mathematical Science, Henan Institute of Science and Technology, Xinxiang, China

²Department of Basic Science, Henan Mechanical and Electrical Engineering College, Xinxiang, China

ABSTRACT

In this paper, by using *new linearization* method we present an optimization algorithm for globally solving a class of multiplicative problems which have a broad application in *computational chemistry*, information technology, and so on. By utilizing characteristic of quadratic function, a series of linear relaxation programming problem of the initial problem can be derived and which can provide a reliable lower bound. By means of the subsequent solutions of a sequence of linear relaxation programming problems, the proposed optimization algorithm converges to the global optimal solution of the initial problem. Numerical experimental results show that the proposed algorithm is feasible and effective.

Keywords: Multiplicative problem, *Optimization* algorithm, New linearization method, Linear relaxation programming, Branch and bound

INTRODUCTION

In this paper, we shall consider the following a class of multiplicative problem:

$$(MP): \begin{cases} \min g(x) = \sum_{j=1}^p (\sum_{i=1}^n c_{ji}x_i + d_j)(\sum_{i=1}^n e_{ji}x_i + \beta_j) \\ s.t. Ax \leq b, x \in S^0 = \{x \in R^n : l^0 \leq x \leq u^0\}, \end{cases}$$

where $A \in R^{m \times n}$, $b \in R^m$, $l^0 = (l_1^0, l_2^0, \dots, l_n^0)^T$, $u^0 = (u_1^0, u_2^0, \dots, u_n^0)^T$.

In last many years, many research have been done on solving the multiplicative problem (MP). On the one hand, this is since this problem (M) exists many important applications in economic planning, engineering designing, financial plan, robust optimization [1-6], and so on. On the other hand, it is because that the problem (MP) usually poses many significant theoretical challenge and computational difficulties, i.e., it is well known that the problem (MP) possesses multiple local optimal solutions which are not globally optimal solution. For instances, when $p = 1$, the problem (MP) is a special case of nonconvex programming problems, which is well known to be NP-hard [7]. Therefore, it has attracted interest of many researchers and practitioners.

Up to today, although many feasible algorithms have proposed for solving the multiplicative problem (MP), but to our knowledge, few algorithm has been still designed for globally solving the multiplicative problem (MP).

Many algorithms have proposed for linear multiplicative programming problem (MP). For example, a large number of quadratic programming methods can be obtained to solve the multiplicative problem (MP) in the literatures [8-13]. When feasible region is a polyhedral set and $p \geq 2$, the branch and bound algorithms, the approximating

algorithms, the outcome space branch and bound approaches, the cutting plane methods, the heuristic methods, the monotonic optimization approaches, the simplicial branch and bound algorithms can be used to solve the problem (MP) in [14-28]. Recently, the authors in [29-32] presented several different feasible and effective algorithms for solving the generalized linear multiplicative programming problem (MP). In addition, some feasible global optimization algorithms for solving generalized nonlinear multiplicative programming problem have been proposed in [33-35].

In this article, we will present a feasible algorithm for the problem (MP) by solving a series of linear relaxation programming problems over partitioned subsets. To globally solve the problem (MP), we first transform the problem (MP) into an equivalent quadratic programming problem (EP), then, a new linearization technique is used to systematically convert the problem (EP) into a sequences of linear relaxation programming problems. The optimal solutions of these transformed problems can approximate sufficiently the global optimal solution of the problem (EP) by a successive partition process. Finally, numerical examples and their computational results are given, and numerical results show that the proposed algorithm can be used to solve all the test problems in computing the global optimal solutions of the multiplicative problem (MP) within a given tolerance condition.

The paper is described as follows. In Sections 2, first we convert the problem (MP) into an equivalent problem (EP), then a new linearizing method is proposed for generating the linear relaxation of the problem (EP). Using the new linearizing method, a branch and bound algorithm is established for globally solving the (EP), and the convergence of the proposed algorithm is proved. Some numerical results are reported in Section 3 and Section 4 provides some concluding remarks.

RESULTS AND DISCUSSION

Firstly, we shall convert the objective function of the problem (MP) into an equivalent quadratic function.

$$\begin{aligned} g(x) &= \sum_{j=1}^p \left(\sum_{i=1}^n c_{ji} x_i + d_j \right) \left(\sum_{i=1}^n e_{ji} x_i + \beta_j \right) \\ &= \sum_{i=1}^n \sum_{k=1}^n \left(\sum_{j=1}^p c_{ji} e_{jk} \right) x_i x_k + \sum_{i=1}^n \left(\sum_{j=1}^p d_j e_{ji} + \beta_j c_{ji} \right) x_i \\ &= \sum_{j=1}^p \sum_{i=1}^n \sum_{k=1}^n c_{ji} e_{jk} x_i x_k + \sum_{j=1}^p \left(d_j \sum_{i=1}^n e_{ji} x_i + \beta_j \sum_{i=1}^n c_{ji} x_i \right) + \sum_{j=1}^p d_j \beta_j \end{aligned}$$

Let

$$Q_{ik} = \sum_{j=1}^p c_{ji} e_{jk}, \quad i = 1, \dots, n, \quad k = 1, \dots, n, \quad B_i = \sum_{j=1}^p (d_j e_{ji} + \beta_j c_{ji}), \quad i = 1, \dots, n, \quad E = \sum_{j=1}^p d_j \beta_j.$$

Define the matrix $Q = (Q_{ik})_{n \times n}$, where Q_{ik} is a component of Q , define the vector B , where B_i is i_{th} a component of B . Obviously, the problem (MP) can be converted into the following equivalent quadratic programming problem:

$$(EP) : \begin{cases} \min g(x) = x^T Q x + B^T x + E \\ s.t. Ax \leq b, \quad x \in S = \{x \in R^n : l \leq x \leq u\}. \end{cases}$$

Let

$$S^k = \{x \in R^n : l^k \leq x \leq u^k\} \subseteq S^0, \quad l^k = (l_1^k, \dots, l_n^k)^T, \quad u^k = (u_1^k, \dots, u_n^k)^T.$$

Set λ_{\min} be the smallest eigen values of Q , and let

$$\theta = \begin{cases} 0, & \text{if } \lambda_{\min} \geq 0; \\ |\lambda_{\min}| + \tau, & \text{if } \lambda_{\min} < 0. \end{cases}$$

where $\tau \geq 0$, then $Q + \theta I$ is a positive semi-definite matrix. Then

$$g(x) = x^T Qx + B^T x + E = x^T (Q + \theta I)x + B^T x - \theta \sum_{j=1}^n x_j^2 + E.$$

According to the characteristic of quadratic function $-x_j^2$ over the interval $[l_j^k, u_j^k]$, we have

$$-x_j^2 \geq 2u_j^k x_j - 2u_j^k l_j^k + (l_j^k)^2, j = 1, \dots, n.$$

$$\begin{aligned} x^T Qx &= (u^k - x)^T (Q + \theta I)(u^k - x)^T - \theta \sum_{j=1}^n x_j^2 + 2(u^k)^T (Q + \theta I)x - (u^k)^T (Q + \theta I)u^k \\ &\geq -\theta \sum_{j=1}^n x_j^2 + 2(u^k)^T (Q + \theta I)x - (u^k)^T (Q + \theta I)u^k \\ &\geq -\theta \sum_{j=1}^n [-2u_j^k x_j + 2u_j^k l_j^k - (l_j^k)^2] + 2(u^k)^T (Q + \theta I)x - (u^k)^T (Q + \theta I)u^k. \end{aligned}$$

Hence, we have

$$g(x) \geq -\theta \sum_{j=1}^n [-2u_j^k x_j + 2u_j^k l_j^k - (l_j^k)^2] + 2(u^k)^T (Q + \theta I)x - (u^k)^T (Q + \theta I)u^k + B^T x + E = g^L(x).$$

Hence, we can establish the linear relaxation programming (RLP) of the problem (EP) over S^k as follows:

$$\text{RLP}(S^k) : \begin{cases} \min g^L(x), \\ \text{s.t. } Ax \leq b, \\ x \in S^k = \{x : l^k \leq x \leq u^k\}. \end{cases}$$

Denote $v(S^k)$ and $LB(S^k)$ as the global optimal value of the problem EP(S^k) and the problem RLP(S^k), respectively. Obviously, we have

$$v(S^k) \geq LB(S^k).$$

Theorem 1. For any $x \in S^k$, we have the following conclusions:

$$\lim_{\|u^k - l^k\| \rightarrow 0} |g(x) - g^L(x)| = 0.$$

Proof. By the expression of the function $g(x)$ and $g^L(x)$, for any $x \in S^k$, we can get that

$$\begin{aligned} g(x) - g^L(x) &= x^T (Q + \theta I)x + d^T x - \theta \sum_{j=1}^n x_j^2 - [-\theta \sum_{j=1}^n [-2u_j^k x_j + 2u_j^k l_j^k - (l_j^k)^2]] \\ &\quad + 2(u^k)^T (Q + \theta I)x - (u^k)^T (Q + \theta I)u^k + d^T x \\ &= x^T (Q + \theta I)x - [2(u^k)^T (Q + \theta I)x - (u^k)^T (Q + \theta I)u^k] \\ &\quad - \theta [(\sum_{j=1}^n (-2u_j^k x_j + 2u_j^k l_j^k - (l_j^k)^2) - \sum_{j=1}^n x_j^2)] \\ &\leq \left\| (u^k - x)^T (Q + \theta I)(u^k - x) + \theta \sum_{j=1}^n [2u_j^k x_j - 2u_j^k l_j^k + (l_j^k)^2 - x_j^2] \right\| \\ &\leq \left\| (u^k - x)^T (Q + \theta I)(u^k - x) + \theta \sum_{j=1}^n (x_j - l_j^k)(u_j^k - x_j + u_j^k - l_j^k) \right\| \\ &\leq \|Q + \theta I\| \|u^k - l^k\|^2 + 2\theta \|(u^k - l^k)^T (u^k - l^k)\| \\ &\leq \|Q + \theta I\| \|u^k - l^k\|^2 + 2\theta \|u^k - l^k\|^2 \end{aligned}$$

Therefore,

$$\lim_{\|u^k - l^k\| \rightarrow 0} |g(x) - g^L(x)| = 0.$$

Based on the above new linearization method, we can construct the linear relaxation programming (LRP) of the problem (EP), which can offer a valid lower bound for the global optimal value of the problem (EP) over rectangle S^k .

Based on the former linearization technique, an effective branch and bound algorithm is proposed for globally solving the problem (EP). To compute the global optimization solution of the problem (EP), the proposed algorithm needs to solve a series of linear relaxation programming problem over partitioned subsets of S^0 . Furthermore, to guarantee that the proposed algorithm is convergent to the global optimal solution.

The proposed algorithm is based on subdividing the set S^0 into two sub-hyper-rectangles, and each sub-hyper-rectangle is corresponding to a node of a branch and bound tree, and each node is corresponding to a linear relaxation programming problem in the associated sub-hyper-rectangle. Therefore, at iteration k of the proposed algorithm, assume that we get a collection of active nodes represented as Ω_k , say, each is associated with a hyper-rectangle $S \subseteq S^0$, $\forall S \in \Omega_k$. For each such node S , we will calculate a lower bound $LB(S)$ of the problem (EP) by solving the problem (RLP). Therefore, the lower bound of global optimal value of the problem (EP) on the whole initial rectangle S^0 at iteration k is given by $LB_k = \min\{LB(S), \forall S \in \Omega_k\}$.

As the optimal solution of the relaxation linear programming problem (RLP) is feasible to the problem (EP), we renew the upper bound UB_k of the problem (EP), if necessary. Therefore, the active nodes collection Ω_k satisfy $LB(S) < UB$, $\forall S \in \Omega_k$, at any stage k . We now select an active node to partition its associated hyperrectangle into two sub-hyper-rectangles as described below, computing the lower bounds for each new node as before. Upon detecting any non-improving nodes, we can get the collection of active nodes for the next iteration, and this process is repeated until the condition of the convergence is satisfied.

The critical element in guaranteeing convergence to a global minimum is the choice of a suitable partitioning strategy. In our paper we choose a simple and standard bisection rule. This method is sufficient to ensure convergence since it drives all the intervals to zero for all variables. This branching rule is as follows.

Suppose that the rectangle $S^k = [l^k, u^k] \subseteq S^0$ will be divided. Then we will choose the branching variable x_p satisfying $p \in \arg \max\{u^k - l^k : i = 1, 2, \dots, N\}$ and subdivide S^k by partitioning the interval $[l_p, u_p]$ into the subintervals $[l_p, (l_p + u_p)/2]$ and $[(l_p + u_p)/2, u_p]$.

Assume that $LB(S^k)$ be the global optimal value of the problem (RLP) over the rectangle S^k and suppose that $x^k = x(S^k)$ be the global optimal solution of the problem (RLP) over the rectangle S^k . The steps of the proposed branch and bound algorithm are given as follows.

Algorithm statement:

Step 0. (Initializing)

Let initial the iteration number $k := 0$, the initial set of the active node $\Omega_0 = \{S^0\}$; the initial upper bound $UB = \infty$, and the initial set of feasible solution $F := \emptyset$.

Compute $LB_0 := LB(S)$ and $x^0 := x(S)$ by solving the problem (RLP) over rectangle $S \subseteq S^0$.

If x^0 is a feasible solution of the problem (EP), we update feasible set F and the upper bound UB , if necessary. If $UB \leq LB_0 + \varepsilon$, where $\varepsilon > 0$ is a given tolerance constant number, then terminate with x^0 be the optimal solution of the problem (EP). Otherwise, continue to the following Step 1.

Step 1. (Bounding) Choose the midpoint x^{mid} of S^k , if x^{mid} is a feasible solution of the problem (EP), then let $F := F \cup \{x^{mid}\}$. And define the upper bound by $UB := \min_{x \in F} g(x)$.

If $F \neq \emptyset$, denote the best known feasible solution by $b := \arg \min_{x \in F} \varphi(x)$.

Step 2. (Subdividing) According to the proposed partitioning rule, choose the branching variable x_p to subdivide the rectangle S^k into two new sub-hyper-rectangles. Denote the set of new partitioned rectangles by \bar{S}^k .

For each rectangle $S \in \bar{S}^k$, compute the lower bound value g^L of $g(x)$ over the hyper-rectangle S . If the lower bounds g^L satisfies $g^L > UB$, then delete the corresponding sub-hyper-rectangle S from \bar{S}^k , i.e. $\bar{S}^k := \bar{S}^k \setminus S$, and skip to next element of \bar{S}^k .

If $\bar{S}^k \neq \emptyset$, compute $LB(S)$ and $x(S)$ by solving the problem (RLP) over the rectangle $S \in \bar{S}^k$. If $LB(S) > UB$, let $\bar{S}^k := \bar{S}^k \setminus S$; otherwise, renew the obtained UB , F and b if possible, as step 1.

Step 3. (Bounding) The remaining partition set is denoted by $\Omega_k := (\Omega_k \setminus S^k) \cup \bar{S}^k$ which can give a new lower bound $LB_k := \inf_{S \in \Omega_k} LB(S)$.

Step 4. (Termination) Detect non-improving nodes by letting

$$\Omega_{k+1} = \Omega_k \setminus \{S : UB - LB(S) \leq \varepsilon, S \in \Omega_k\}.$$

If $\Omega_{k+1} = \emptyset$, then algorithm terminates with UB be the optimal value of the problem (EP), and b is the global optimal solution. Otherwise, $k := k + 1$, and choose an active node S^k such that $S^k = \arg \min_{S \in \Omega_k} LB(S)$, $x^k := x(S^k)$, and continue to Step 1.

Theorem 2 (convergence theorem) The proposed branch and bound algorithm either stops finitely with the global optimal solution of the problem (EP), or produces an infinite iteration sequence which satisfies any limitation point of the sequence $\{x^k\}$ will be the global solution of the problem (EP) along any infinite branch of the branch-and-bound tree.

EXPERIMENTAL SECTION

To verify the effectiveness and feasibility of the proposed branch and bound algorithm, several test examples are implemented on Intel(R) Core(TM)2 Duo CPU (1.58GHZ) microcomputer, the proposed branch and bound algorithm is coded in C++ procedure, the simplex algorithm is used to solve linear relaxation programming problem and the convergence tolerance is set as $\varepsilon=10^{-6}$. These test examples and their numerical results are given as follows.

Example 1.

$$\begin{cases} \min 3x_1^2 + 2x_2^2 \\ \text{s.t. } x_1 + 2x_2 \leq 4, x_1 - 3x_2 \leq 10, \\ 1 \leq x_1, x_2 \leq 3. \end{cases}$$

Using the proposed algorithm in this paper, the global ε -optimal solution (1.0,1.0) and global ε -optimal value 5.0 is obtained.

Example 2.

$$\begin{cases} \min (x_1 + x_2)(x_1 - x_2) + (x_1 + x_2 + 1)(x_1 - x_2 + 1) \\ \text{s.t. } x_1 + 2x_2 \leq 10, \quad x_1 - 3x_2 \leq 20, \\ 1 \leq x_1, x_2 \leq 3, \end{cases}$$

Using the proposed algorithm in this paper, the global ε -optimal solution (1.0,3.0) and global ε -optimal value -13.0 is obtained.

Example 3.

$$\begin{cases} \min (x_1 + x_2)(x_1 - x_2) + (x_1 + x_2 + 2)(x_1 - x_2 + 2) \\ \text{s.t. } x_1 + 2x_2 \leq 20, \quad x_1 - 3x_2 \leq 20, \\ 1 \leq x_1, x_2 \leq 4, \end{cases}$$

Using the proposed algorithm in this paper, the global ε -optimal solution (1.0,4.0) and global ε -optimal value -22.0 is obtained.

From the numerical results for test examples 1-3, we can get that the proposed branch and bound algorithm is competitive and can be used to globally solve the problem (MP).

CONCLUSION

In this article, a branch and bound algorithm is proposed to solve the problem (MP). In the algorithm, a new linearization method is proposed. By utilizing the method the problem (MP) can be transformed into a series of linear relaxation programming problems, which can be used to compute the lower bound of the optimal value of the problem (MP). The proposed algorithm is convergent to the global optimal solution of the problem (MP) by solving a series of linear relaxation programming problems. Numerical results show the feasibility of the proposed algorithm.

Acknowledgements

This paper is supported by the Foundation for University Key Teacher by the Ministry of Education of Henan Province (2010GGJS-140) and the Natural Science Foundation of Henan Educational Committee (14A110024). The authors acknowledged the help of Prof. Fenghui Wang from Xi'an Jiaotong University while preparing the paper.

REFERENCES

- [1] H. Konno, H. Watanabe. *J. Oper. Soc. Japan*, **1994**, 39: 295-306.
- [2] I. Quesada, I. E. Grossmann. Alternative bounding approximations for the global optimization of various engineering design problems. In I.E. Grossmann, (ed.), *Global Optimization in Engineering Design, Nonconvex Optimization and Its Applications*, Kluwer Academic Publishers, Norwell, **1996**, MA, 9 pp.309-331.
- [3] C. D. Maranas, I. P. Androulakis, C. A. Floudas, A. J. Berger and J. M. Mulvey. *J. Economic Dynamics & Control*, **1997**, 21: 1405-1425.
- [4] J. M. Henderson, R.E. Quandt. *Microeconomic Theory*. **1971**, 2nd edition, McGraw-Hill, New York.
- [5] J. M. Mulvey, R. J. Vanderbei and S. A. Zenios. *Oper. Res.*, **1995**, 43: 264-281.
- [6] T. Matsui, *J. Global Optim.*, **1996**, 9: 113-119.
- [7] H. Konno, T. Kuno. *Math. Program.*, **1992**, 56: 51-64.
- [8] H. Swarup, *Cahier du Centre d'Etudes de Recherche Operationelle*, **1966**, 8: 223-234.
- [9] H.-M. Li, K.-C. Zhang. *Appl. Math. Comput.*, **2006**, 173(1): 394-403.
- [10] H. Wu, K. Zhang. *Appl. Math. Comput.*, **2008**, 197(2): 810-818.
- [11] S.-T. Liu, R.-T. Wang. *Appl. Math. Comput.*, **2007**, 189(2): 1274-1281.
- [12] P. Shen, M. Gu. *Appl. Math. Comput.*, **2008a**, 198(1): 1-11.
- [13] P. Shen, Y. Duan, Y. Ma. *Appl. Math. Comput.*, **2008b**, 201(1-2): 514-526.
- [14] H. Konno, K. Fukaiishi. *A J. Global Optim.*, **2000**, 18: 283-299.
- [15] H. S. Ryoo, N. V. Sahinidis. *J. Global Optim.*, **2003**, 26: 387-418.
- [16] C. Xue, H. Jiao, et al, *J. Henan Normal University*, **2008**, 36 (5): 13-15.
- [17] H. Tuy, N. D. Nghia. *VN J. Math.*, **2003**, 31: 391-402.
- [18] S. Schaible, C. Sadini. *J. Optim. Theory Appl.*, **1995**, 87(2): 441-455.
- [19] Y. Gao, C. Xu, Yongjian Yang. *Appl. Math. Comput.*, **2006**, 179(2): 494-505.
- [20] R. M. Oliveira, P. A. V. Ferreira. *J. Global Optim.*, **2010**, 47: 107-118.

-
- [21] A. M. Ashtiani, P. A. V. Ferreira. *Anais do CNMAC*, **2010**, 3: 377-383.
- [22] H. P. Benson, G.M. Boger. *J. Optim. Theory Appl.*, **2000**, 104(2): 301-322.
- [23] X. J. Liu, T. Umegaki, and Y. Yamamoto. *J. Global Optim.*, **1999**, 4(15): 433-447.
- [24] N.T.H. Phuong, H. Tuy. *J. Global Optim.*, **2003**, 26: 229-259.
- [25] Y. Chen, H. Jiao. *Comput. Oper. Res.*, **2009**, 36: 2573-2579.
- [26] C.-F. Wang, L. San-Yang, S. Pei-Ping. *Appl. Math. Model.*, **2009**, 36(6): 2446-2451.
- [27] T. Kuno, H. Konno. *J. Global Optim.*, **1991**, 1, pp. 267-286, 1991.
- [28] P. Shen, H. Jiao. *Appl. Math. Comput.*, **2006**, 183(1): 328-336.
- [29] C.-F. Wang, S.-Y. Liu. *Comput. Oper. Res.*, **2011**, 38: 1008-1013.
- [30] H. Jiao, Y.R. Guo, P. Shen. *Appl. Math. Comput.*, **2006**, 183(2): 717-728.
- [31] H. Jiao, *Nonlinear Anal.-Theor.*, **2009**, 70: 1113-1123.
- [32] P. Shen, X. Bai, W. Li. *Nonlinear Anal.-Theor.*, **2009**, 71(7-8): 2866-2876.
- [33] H. P. Benson, *J. Optim. Theory Appl.*, **2008**, 137: 105-120.
- [34] H. Konno, T. Kuno, and Y. Yajima. *J. Global Optim.*, **1994**, 4: 47-62.
- [35] B. Jaumard, C. Meyer, H. Tuy. *J. Global Optim.*, **1997**, 10: 229-256.