



## ***Antioxidants: A Review***

<sup>1</sup>Sulekha Mandal, <sup>1</sup>Satish Yadav, <sup>2</sup>Sunita Yadav, <sup>3\*</sup>Rajesh Kumar Nema

<sup>1</sup>Mahatma Gandhi College of Pharmaceutical Sciences, Jaipur

<sup>2</sup>Dept. of Botany, University of Rajasthan, Jaipur

<sup>3</sup>Rishiraj College of Pharmacy, Indore, India

---

### **Abstract**

Antioxidants are an inhibitor of the process of oxidation, even at relatively small concentration and thus have diverse physiological role in the body. Antioxidant constituents of the plant material act as radical scavengers, and helps in converting the radicals to less reactive species. A variety of free radical scavenging antioxidants is found in dietary sources like fruits, vegetables and tea, etc. This review presents some information about the antioxidant/antiradicals and their role in our body and also their presence in spices and herbs.

**Key Words:** Antioxidants, Aging, Free radicals, Medicinal plants, Spices, Herbs and Vitamins.

---

### **Introduction**

Oxygen is absolutely essential for the life of aerobic organism but it may become toxic if supplied at higher concentrations. Dioxygen in its ground state is relatively unreactive; its partial reduction gives rise to active oxygen species (AOS) such as singlet oxygen, super oxide radical anion, hydrogen peroxide etc. This is partly due to the oxidative stress that is basically the adverse effect of oxidant on physiological function. Free oxygen radicals plays cardinal role in the etiology of several diseases like arthritis, cancer, atherosclerosis etc. The oxidative damage to DNA may play vital role in aging [1]and the presence of intracellular oxygen also can be responsible to initiate a chain of inadvertent reaction at the cellular level and these reaction cause damage to critical cell biomolecules. These radicals are highly toxic and thus generate oxidative stress in plants. Plants and other organism have in built wide range of mechanism to combat with these Free Radical problems. Free radicals are an atom or molecule that bears an unpaired electron and is extremely reactive, capable of engaging in rapid change reaction that destabilize other molecules and generate many more free radicals. In plants and animals these free radicals are deactivated by *antioxidants*. These antioxidants act as an inhibitor of the process of oxidation, even at relatively small concentration and thus have diverse physiological role in the

body. Antioxidant constituents of plant materials act as radical scavengers, and convert the radicals to less reactive species.

Plants have developed an array of defense strategies (antioxidant system) to cope up with oxidative stress. The antioxidative system includes both enzymatic and non-enzymatic systems. The non enzymatic system includes ascorbic acid (vitamin C);  $\alpha$ -tocopherol, carotenes etc. and enzymic system include superoxide dismutase (SOD), catalase (CAT), peroxidase (POX), ascorbate peroxidase (APX), glutathione reductase (GR) and polyphenol oxidase (PPO) etc. The function of this antioxidant system is to scavenge the toxic radicals produced during oxidative stress and thus help the plants to survive through such conditions.

Spices and herbs in food as medicine is a current hot trend that is capturing everyone's imagination with images of a new *magic bullet or fountain of youth*. The intake of antioxidant compounds present in food is an important health-protecting factor. Natural antioxidants present in foods and other biological materials have attracted considerable interest because of their presumed safety and potential nutritional and therapeutic effects. Because extensive and expensive testing of food additives is required to meet safety standards, synthetic antioxidants have generally been eliminated from many food applications. The increasing interest in the search for natural replacements for synthetic antioxidants has led to the antioxidant evaluation of a number of plant sources.

Antioxidants that have traditionally been used to inhibit oxidation in foods also quench dreaded free radicals and stop oxidation chains *in-vivo*, so they have become viewed by many as nature's answer to environmental and physiological stress, aging, atherosclerosis, and cancer. The nutraceutical trend towards doubling the impact of natural antioxidants that stabilize food and maximize health impact presents distinct challenges in evaluating antioxidant activity of purified individual compounds, mixed extracts, and endogenous food matrices and optimizing applications.

It is well known that Mediterranean diet, which is rich in natural antioxidants, leads to a limited incidence of cardio- and cerebrovascular diseases[2]. It is known that compounds belonging to several classes of phytochemical components such as phenols, flavonoids, and carotenoids are able to scavenge free radical such as  $O_2^{\cdot -}$ ,  $OH^{\cdot}$ , or lipid peroxy radical  $LOO^{\cdot}$  in plasma [3]. The effective intake of single food antioxidants and their fate in the human body have been defined only for a few compounds [4-5]. It is reasonable that the higher the antioxidant content in foods is, the higher the intake by the human body will be.

Natural antioxidants occur in all parts of plants. These antioxidants include carotenoids, vitamins, phenols, flavonoids, dietary glutathione, and endogenous metabolites [6]. Plant-derived antioxidants have been shown to function as singlet and triplet oxygen quenchers, free radical scavengers, peroxide decomposers, enzyme inhibitors, and synergists[7]. The most current research on antioxidant action focuses on phenolic compounds such as flavonoids. Fruits and vegetables contain different antioxidant compounds, such as vitamin C, vitamin E and carotenoids, whose activities have been established in recent years. Flavonoids, tannins and other phenolic constituents

Present in food of plant origin are also potential antioxidants[8-9]. These components include:

- Nutrient-derived antioxidants like ascorbic acid (vitamin C), tocopherol and tocotrienols (vitamin E), carotenoids, and other low molecular weight compounds such as glutathione and lipoic acid.

- Antioxidant enzymes, e.g., super oxide dismutase, glutathione peroxidase, and glutathione reductase, which catalyze free radical quenching reactions.
- Metal binding proteins, such as ferritin, lactoferrin, albumin, and ceruloplasmin that sequester free iron and copper ions that are capable of catalyzing oxidative reactions.
- Numerous other antioxidant phytonutrients present in a wide variety of plant foods.

### Conclusion

The most important free radical in biological systems is radical derivatives of oxygen with the increasing acceptance of free radical as commonplace and important biochemical intermediate. Antioxidants are believed to play a very important role in the body defense system against reactive oxygen species (ROS), which are the harmful byproducts generated during normal cell aerobic respiration [10]. Increasing intake of dietary antioxidants may help to maintain an adequate antioxidant status and, therefore, the normal physiological function of a living system [11]. To protect the cells and organ systems of the body against reactive oxygen species, humans have evolved a highly sophisticated and complex antioxidant protection system. It involves a variety of components, both endogenous and exogenous in origin, that function interactively and synergistically to neutralize free radicals.

### References

1. Cheeseman K.H. and Slater T.F. (1993) Free radicals in medicine. Churchill Livingstone Pub. British Med. Bull., 49: 479-724.
2. Frei B., Stocker R. and Ames B. N. (1988) *Proc. Natl. Acad. Sci. U.S.A.*, 85: 9748-9752.
3. Gutteridge J. M. C. and Halliwell B. (2000) *Ann. N. Y. Acad. Sci.*, 899: 136-147.
4. Hertog M. G. L., Feskens E. J. M., Hollman P. C. H., Katan M. B. and Krombhout D.. *Lancet*, 342: 1007-1011.
5. Kaur C. and Kapoor H. C. (2001) *Int. J. Food Sci. Technol.*, 36: 703-725.
6. Larson R. A. (1988) *Phytochemistry*, 4: 969-978.
7. Manach C., Morand C., Crespy V., Demigne C., Texier O., Regeat F. and Remesy C. (1998) *FEBS Lett.*, 426: 331-336.
8. Ramarathnam N., Osawa T., Ochi H. and Kawakishi S. (1995) *Trends Food Sci. Technol.*, 6: 7582.
9. Record I. R., Dreosti I. E. and McInerney J. K. (2001), *Brit. J. Nutr.*, 85: 459-464.
10. Salah N., Miller N. J., Paganga G., Tijburg L., Bolwell G. P. and Rice- Evans C. (1995), 322(2): 339-346.
11. Van Acker S. A. B. E., Van den Vijgh W. J. F. and Bast F. (1996), 20(3): 331-342.