



Determination of sugars by acid hydrolysis of peanut husk (*Arachis hypogaea*) by standard methods

Santhosh Kumar S.^{*1}, Sudarshan S.² and Venkatesh R.²

¹Department of Studies in Biochemistry, Shivagangothri, University of Davanagere, Davangere, Karnataka, India

²Department of Biochemistry, Yuvaraja's College, University of Mysore, Mysore, Karnataka, India

ABSTRACT

Peanut husk is a carbohydrate rich source that can be converted to reducing sugars. Peanut husk was hydrolyzed using (0.2N) sulphuric acid at a temperature 60-65°C. It was observed that the degradation has significant effect with respect to amount of husk taken and in turn sugar yield is around 40-50% each which is estimated by Bertrand's, Lane-Eynon and Benedict's Method.

Keywords: Degradation, hydrolysis, peanut husk, sugar, estimation.

INTRODUCTION

Peanut is drought tolerant and resistant to water logging [1] and grows in various soil conditions [2]. Peanut husk is the outer most layers of peanut seed, also called peanut hull. Peanut husk is an agricultural waste material obtained from milling of the peanut seeds and constitutes about 20% of 34.4 million metric tons of peanut seeds produced annually in the world [3, 4].

Peanut husk predominantly contains hemicelluloses (approx.15.2%), and cellulose (approx.18.5%) the lignocelluloses biomass are hydrolyzed to convert hemi cellulose and cellulose into sugar [5]. According to Badger [6] there are two types of hydrolysis i.e. enzymatic and chemical hydrolysis. Chemical hydrolysis was selected because it is relatively low cost and fast [7, 8].The dilute acid hydrolysis of lignocelluloses biomass was run with operating condition of Sulphuric acid (0.2 N) Concentration, 60-65°C, at various amount of peanut husk.

The standard methods adopted for estimation are;

(i) Bertrand's method [9] is based on the reducing action of sugar on the alkaline solution of tartarate complex with cupric ion; the cuprous oxide formed is dissolved in warm acid solution of ferric alum. The ferric alum is reduced to FeSO₄ which is titrated against standardized KMnO₄; Cu equivalence is correlated with the table to get the amount of reducing sugar.

(ii) In Lane-Eynon method [10] sugar solution is taken in the burette and known volume of Fehling solution is taken in conical flask. This is titrated at a temperature 60-65°C. Titration is continued till it acquires a very faint blue color; add 3 drops of methylene blue indicator. The dye is reduced to a colorless compound immediately and the color changes from blue to red (at the end point).

(iii) Benedict quantitative reagent gives a visual clear end point which turn blue to white by using potassium thiocyanate which converts the red cuprous oxide to white crystals of cuprous thiocyanate, it helps in visual view.[11]

EXPERIMENTAL SECTION

The hydrolysis of peanut husk was carried out at constant stirring using 50 ml of 0.2N sulphuric acid temperature in a hotplate, equipped with a temperature controller, and continuously shaken during the operation. Initially, 50mL of 0.2 N sulphuric acid solution 20 mesh peanut husk were put into the beaker and kept under hot plate as well as the temperature controller was adjusted such that the temperature of the mixture is about 60-65°C. The reaction was expected to be at constant temperature (isothermal), but before that temperature was achieved, reaction has occurred. The hydrolyzate was neutralized to bring the pH to 7 by the addition of calcium carbonate and activated carbon, followed by filtration. The concentration of reducing sugar was analyzed by Benedict's Bertrand's and Lane-Eynon standard procedures.

RESULTS AND DISCUSSION

By varying the amount of peanut husk 1, 2, 3, 4 and 5g respectively at constant temperature (60-65°C) and concentration of sulphuric acid is 0.2N is fixed constant. The experiment resulted in the data of reducing sugar concentrations at 3 hour were reported below **Table 1** and there corresponding data are plotted which are shown in **figure 1, 2 and 3** respectively.

Table 1: Amount of Reducing Sugar Estimated by different methods

Weight of peanut husk	Sugar Estimation Benedict's method(g)	Sugar Estimation by Bertrand's Method(g)	Sugar Estimation by Lane-Eynon Method(g)
1.012	0.232	0.235	0.233
2.009	0.464	0.470	0.466
3.019	0.696	0.705	0.699
4.002	0.931	0.939	0.929
5.001	1.162	1.173	1.160




Figure 1: Estimation of reducing sugar by Benedict's method




Figure 2: Estimation of reducing sugar by Bertrand's method




Figure 3: Estimation of reducing sugar by Lane-Eynon's method

- Amount of peanut husk taken (gm)
- Amount of reducing sugar estimated (gm)

CONCLUSION

While there are some uses generally, peanut husk is used as fuel but it is still often considered a waste product in the mill and therefore often either burned in the open or dumped on wasteland. In the present work, we have applied simple hydrolysis process to obtain reducing sugars which is very good consumable source of energy and the yield percent also runs up to 40-50% which is authentically reported by analytical standard procedures in an economical way.

Acknowledgement

The author would like to thank all professors in the department of bio-chemistry, Shivagangothri, University of Davanagere, Davanagere, for their support and encouragement in carrying out of the work.

REFERENCES

- [1] H Doggett, Sorghum, *Longman scientific and Technical, London (1998)*, 2, 1-3.

-
- [2] SL Dillon, FM Shapter, RJ Henry, G Cordeiro, L Izquierdo, LS Lee, Domestication to crop improvement; *Genetic Resources for sorghum and Saccharum (Andropogoneae)*. *Annals of Botany* **2007**, 5, 975-989
- [3] USDA foreign agricultural service.
- [4] Board on science and technology for international development, office of international affairs, National Research Council of the National Academies, *Lost Crops of Africa; vol; Grains*, The National Academy Press, Washington, DC **1996**,
- [5] Hahn-Hagerdal, B Galbe, FM Gorwa-Grausland, Liden, G Zacchi, *Science Direct, Elsevier*, **2006**.
- [6] PC Badger, *Trends in New Crops and New Uses*, ASHS Press., Alexandria VA, **2002**, 17-21.
- [7] E Palmqvist, H, Hagerdal, *Bioresource Technology, Elsevier*, **2000**, 74, 25-33.
- [8] MJ Taherzadeh, R Eklund, L Gustafsson, C Niklasson, G Liden, *Ind. Eng. Chem. Res., American Chemical Society*, **1997**, 36, 4659-4665.
- [9] <http://www.raktuo-kasei.net/textile/8.index.html>
- [10] Sausen Silmi, HS Canarsie, Brooklyn, *Summer Research Programme for Teachers, Determining the Concentration of Solution: Beer's Law* **1997**
- [11] GRF Davis, *Clinica Chimica Acta*, **1963**, 8, 635-636.