



ISSN No: 0975-7384  
CODEN(USA): JCPRC5

*J. Chem. Pharm. Res.*, 2011, 3(5):407-419

---

## **DNA Binding and Cleavage Studies of Fe(II) and Zn(II) Complexes containing Mixed Ligand of 1,10-phenanthroline and 2-hydroxy-4-methyl-1,8-naphthyridine**

**B. Sreekanth<sup>a</sup>, G. Krishnamurthy<sup>b\*</sup>, H. S. Bhojya Naik<sup>c</sup>, T. K. Vishnuvardhan<sup>d</sup>, N. D. Shashikumar<sup>b</sup> and M. R. Lokesh<sup>b</sup>**

<sup>a</sup>Department of Biotechnology, Acharya Institute of Technology, Bangalore, Karnataka, INDIA

<sup>b</sup>Department of Chemistry, Sahyadri Science College, Shimoga, Karnataka, INDIA

<sup>c</sup>Department of PG Studies and Research in Industrial Chemistry, School of Chemical Sciences, Kuvempu University, Shankaraghatta, Shimoga, Karnataka

<sup>d</sup>Department of Chemistry, Acharya Institute of Technology, Bangalore, Karnataka, INDIA

---

### **ABSTRACT**

The DNA binding ability of the novel complexes  $[Fe(L_2)_2L_1] (PF_6)_2$  [Complex (1)] and  $[Zn(L_2)_2L_1] (PF_6)_2$  [Complex (2)] containing bioactive mixed ligand of the type  $L_1=2$ -hydroxy-4-methyl-1,8-naphthyridine and  $L_2 =1,10$ -phenanthroline were synthesized and structurally characterized by elemental analysis, IR and  $^1H$  NMR spectral studies. The intrinsic binding constant  $K_b$  has been estimated at room temperature. The binding constant of  $1.8 \times 10^4 M^{-1}$  for complex (1) and  $2.1 \times 10^4 M^{-1}$  for complex (2) in 5 mM Tris-HCl/50 mM NaCl buffer at pH 7.2, respectively, as obtained from absorption spectra indicate that the complexes intercalate between the base pairs of the CT-DNA tightly. The oxidative cleavage activity of the complexes (1) and (2) were studied by using gel electrophoresis and the results show that complexes have potent nuclease activity.

**Key words:** Fe(II) and Zn(II) Complexes, DNA Binding, Cleavage studies.

---

### **INTRODUCTION**

The interactions of metal complexes with DNA(Deoxyribonucleic acid) is a recent focus of research interest in bioinorganic chemistry. The binding and cleavage of DNA by metal

complexes and it is related to the utility of such metal complexes in the design and the development of synthetic restriction enzymes, new drugs DNA foot printing agents etc.[1-5]. Metal complexes have been found to be particularly useful for the above mentioned purposes because of their potential to bind DNA via multitude of interactions and to cleave the duplex [6-8] by virtue of their intrinsic chemical, electrochemical and photochemical reactivities [9-11]. Indeed, there is already a considerable literature involving the practical use of transition metal complexes as chemical nucleases [12-24]. However, most of these complexes contain only planar aromatic ligands and investigations of such complexes with ligands containing substituents as DNA-binding reagents have been relatively few. In fact, some of these complexes also exhibit interesting properties like anti-inflammatory, anticancer and antitumor activities upon binding to DNA [25-29]. The features common to these complexes are that the molecule has a high affinity for double-stranded DNA, and that the molecule also binds a redox active metal ion cofactor. The ligands or the metal in these complexes can be varied in an easily controlled manner to facilitate an individual application [30]. All the studies reveal that modification of the metal or ligands would lead to subtle or substantial changes in the binding modes, location and affinity giving rise to changes to explore various valuable conformation or site-specific DNA probes and potential chemotherapeutic agents [31-34].

Studies of mixed ligand transition metal complexes, which bind at specific sites along a DNA strand as reactive models for protein-nucleic acid interaction, provide routes toward rational drug design as well as means to develop sensitive chemical probes for DNA. Thus, a number of metal chelates are of current interest for important applications in nucleic acid chemistry as probes of DNA structure in solution, reagents for mediation of strand scission of duplex DNA under physiological conditions, and chemotherapeutic agents and in genomic research [35-37].

There have been immense interest to develop the virtually unexplored chemistry of non porphyrinic 3d-metal complexes containing bio-essential metal ion like iron considering the utility of such complexes as viable substitute of porphyrinic compounds in PDT(Photodynamic therapy) applications [38, 39].

The role of zinc in a wide range of cellular processes, including cell proliferation, reproduction, immune function and defense against free radicals, has been well established [40]. Zinc is considered the most abundant trace intracellular element, and there exists increasing evidence that zinc plays an important role in both genetic stability and function [41]. In vitro, significant amounts of zinc are incorporated in the nuclei [42]. It is clear that, mechanistically, zinc has a significant impact on DNA as a component of chromatin structure, DNA replication and transcription and DNA repair [43]. Zinc is a component of more than 3000 zinc-associated transcription factors, including DNA-binding proteins with zinc fingers, and more than 300 enzymes, including copper/zinc superoxide dismutase (CuZnSOD) and several proteins involved in DNA repair [44-46]. Thus, zinc plays an important role in protecting cellular components from oxidation and damage to DNA [47].

Although DNA interactions of number of mixed ligand complexes previously appeared [48-53], there is still scope to design and Schiff base containing 1,10-phenantroline/2-hydroxy-4-methyl-1,8-naphthyridine with the Fe(II) and Zn(II) as new chemical nucleases. Bearing these facts in

mind, the nuclease activity of mixed-ligand complexes of Fe(II) and Zn(II) containing 1,10-phenanthroline/2-hydroxy-4-methyl-1,8-naphthyridine is reported herein.

## EXPERIMENTAL SECTION

The reagents and solvents used were of AR grade. Solvents were purified and used. Zinc acetate, ferrous sulphate,  $\text{NH}_4\text{PF}_6$  (ammonium hexafluorophosphate), DMSO (Dimethylsulphoxide) Tris-HCl buffer were purchased from qualigens (Mumbai, India). Ligand 1,10-phenanthroline was purchased from Sigma Aldrich(Bangalore). CT-DNA (Calf thymus Deoxyribonucleic acid) and pUC 19 DNA(plasmid University of California 19 Deoxyribonucleic acid) were purchased from Bangalore Gene, Bangalore, India.

### Synthesis of Ligand (2-hydroxy-4-methyl-1,8-naphthyridine)

A mixture of 2-aminopyridine (1 g, 0.01 mol) and ethylacetoacetate (1.3 g, 0.013 mol) was irradiated in a microwave oven for 8 min. and 2 to 4 drops of  $\text{H}_2\text{SO}_4$  was added and again irradiated for 3 min. A yellowish product precipitated out with a yield of 82%. It was washed with cold ethanol, dried under vacuum and recrystallised from ethylacetate. Anal. Calcd for  $\text{C}_9\text{H}_8\text{N}_2\text{O}$ . calcd. (%) : C 67.50; H 5.00; N 8.75; O 10.00. Found (%): C 67.48; H 4.98; N 8.68; O 9.97. IR, KBr pellets ( $\nu$ ,  $\text{cm}^{-1}$ ): 1462 ( $\nu(\text{C}=\text{N})$ ); 3149 ( $\nu(\text{C}-\text{H}, \text{Ar}-\text{H})$ ); 3258 ( $\nu(\text{OH})$ ).  $^1\text{H}$  NMR ( $\delta$ , ppm): 2.60 (m, 3H,  $\text{CH}_3$ ), 4.70(s, H, Ar-OH), 7.80(m, 4H, Ar-H), 8.8(m, 8H, ArN-H), 8.1(m, 2H,  $\text{CH}=\text{N}$ ).

### Synthesis of Metal Complexes

#### Synthesis of Metal Complex $[\text{Fe}(\text{phen})_2\text{SO}_4]3\text{H}_2\text{O}$

Ferrous sulphate (0.278g, 1 mmol) and 1,10-phenanthroline(0.396g, 1 mmol) were dissolved in hot methanolic solution and refluxed on the water bath for 3h. The contents were cooled to obtain precipitate. The complex was filtered and dried under vacuum before being recrystallized in acetone. Yield of the complex 80%.

#### Synthesis of $[\text{Fe}(\text{L}_2)_2\text{L}_1] (\text{PF}_6)_2$ [Complex (1)]

50 ml of hot methanolic solution of  $[\text{Fe}(\text{phen})_2\text{SO}_4]3\text{H}_2\text{O}$  (0.56 g, 1 mmol) was added to an hot methanolic solution of 2-hydroxy-4-methyl-1,8-naphthyridine( $\text{L}_1$ ) (0.161g, 1mmol). The mixture was refluxed on the water bath for 3h. The contents were cooled and precipitated by the addition of hot methanolic solution of  $\text{NH}_4\text{PF}_6$  to the filtrate. The complex  $[\text{Fe}(\text{L}_2)_2\text{L}_1] (\text{PF}_6)_2$  was filtered and dried under vacuum before being recrystallized in acetone. Yield of the complex 78%. Anal. Calcd for  $\text{C}_{33}\text{H}_{24}\text{N}_6\text{OP}_2\text{F}_{12}\text{Fe}$ : C, 43.69; H, 2.69; N, 9.35; O, 1.86; Fe, 6.14. Found: C, 43.65; H, 2.67; N, 9.29; O, 1.81; Fe, 6.09. IR, KBr pellets ( $\text{cm}^{-1}$ ): 721(C-H, out of plane bend), 1636( $\nu(\text{C}=\text{N})$ ), 3051( $\nu(\text{C}-\text{H})$ ), 3367( $\nu(\text{O}-\text{H})$ ).  $^1\text{H}$  NMR ( $\delta$ , ppm): 2.40 (m, 3H,  $\text{CH}_3$ ), 4.80(s, H, Ar-OH), 8.30(m, 4H, Ar-H), 8.70(m, 8H, ArN-H), 7.80(m, 2H,  $\text{CH}=\text{N}$ ).

#### Synthesis of Metal Complex $[\text{Zn}(\text{phen})_2](\text{CH}_3\text{COO})_2.3\text{H}_2\text{O}$

Zinc acetate(0.44 g, 1 mmol) and 1,10-phenanthroline(0.396g, 1 mmol) were dissolved in hot methanolic solution and refluxed on the water bath for 3h. The contents were cooled to obtain precipitate. The complex was filtered and dried under vacuum before being recrystallized in acetone. Yield of the complex 75%.

**Synthesis of  $[\text{Zn}(\text{L}_2)_2\text{L}_1] (\text{PF}_6)_2$  [Complex (2)]**

To a hot methanolic solution of 2-hydroxy-4-methyl-1,8-naphthyridine( $\text{L}_1$ )(0.161g, 1mmol) 50ml of hot methanolic solution of  $[\text{Zn}(\text{phen})_2] (\text{CH}_3\text{COO})_2 \cdot 3\text{H}_2\text{O}$  (0.58 g, 1 mmol) was added. The mixture was refluxed on the water bath for 3h. The contents were cooled and precipitated by the addition of hot methanolic solution of  $\text{NH}_4\text{PF}_6$  to the filtrate. The complex  $[\text{Zn}(\text{L}_2)_2\text{L}_1] (\text{PF}_6)_2$  was filtered and dried under vacuum before being recrystallized in acetone. Yield of the complex 74%. Anal. calcd for  $\text{C}_{33}\text{H}_{24}\text{N}_6\text{OP}_2\text{F}_{12}\text{Zn}$ : C, 43.24; H, 2.63; N, 9.18; O, 1.75; Zn, 6.98. Found: C, 43.28; H, 2.68; N, 9.21; O, 1.78; Zn, 6.99. IR, KBr pellets ( $\text{cm}^{-1}$ ): 722(C-H, out of plane bend), 1554v(C=N), 3067v(C-H), 3229v(O-H).  $^1\text{H}$  NMR ( $\delta$ , ppm): 2.40 (m, 3H,  $\text{CH}_3$ ), 4.80(s, H, Ar-OH), 8.30(m, 4H, Ar-H), 8.90(m, 8H, ArN-H), 8.10(m, 2H, CH=N).

The structure of the investigated complexes is given in Fig. 1.




FIGURE 1: The structure of the complexes where M is  $\text{Fe}^{2+}$  or  $\text{Zn}^{2+}$

**Spectral Measurements**

Melting points were determined in open capillaries and are uncorrected. Microanalysis (C, H, and N) were performed in Carlo-Erba 1106 model 240 Perkin-Elmer analyzer at IISc Bangalore. The molar conductivity in Dimethylformamide(DMF) ( $10^{-3}\text{M}$ ) at room temperature were measured using Equiptronics digital conductivity meter. IR spectra were recorded with Shimadzu model FT-IR spectrophotometer by using KBr pellets at Acharya Pharmacy College, Bangalore. Bruker FT-NMR Spectrophotometer (400 MHz) was used for recording  $^1\text{H}$ -NMR spectra at  $25^\circ\text{C}$  in MeOD (detoriated methanol) with TMS(tetra methyl silane) as the internal reference at IISc, Bangalore. UV-visible absorption spectra were recorded at department of Industrial Chemistry, Kuvempu University, Shankaraghatta, using Shimadzu model UV-1650PC spectrophotometer at room temperature. Viscosity measurements were carried out on semi-micro dilution capillary viscometer (Viscomatic Fica MgW) with a thermostated bath D40S at room temperature. Thermal denaturation studies were carried out with a Perkin-Elmer Lambda 35 Spectrophotometer.

**DNA Binding and Cleavage experiments**

The concentration of CT-DNA per nucleotide  $[\text{C}(\text{p})]$  was measured by using its known extinction coefficient at 260 nm ( $6600 \text{ M}^{-1} \text{ cm}^{-1}$ ) [54]. Tris HCl-buffer [5mM tris(hydroxymethyl) amino methane, pH 7.2, 50 mM NaCl] was used for the absorption, viscosity and thermal denaturation experiments.

Absorption titration experiments were carried out by varying the DNA concentration (0-100  $\mu\text{M}$ ) and maintaining the metal complex concentration constant. Absorption spectra were recorded after each successive addition of DNA and equilibration (approximately 10 minutes). The absorption data were analyzed for an evaluation of the intrinsic binding constant  $K_b$  using reported procedure [55].

Each viscosity experiment was performed three times and average flow time was calculated. Data were presented as  $(\eta/\eta_0)$  versus binding ratio where  $\eta$  is viscosity of DNA in the presence of complex and  $\eta_0$  is the viscosity of DNA alone.

Thermal denaturation experiments were carried out with a Shimadzu Model UV-160A spectrophotometer coupled to a temperature controller (Model TCC-240A) by monitoring the absorption of CT-DNA (50  $\mu\text{M}$ ) at 260 nm at various temperatures, both in the presence (5-10  $\mu\text{M}$ ) and absence of each complex. The melting temperature ( $T_m$ , the temperature at which 50% of double stranded DNA becomes single stranded) and the curve width ( $\sigma_T$ , the temperature range between which 10% and 90% of the absorption increases occurred) were calculated as reported [56, 57].

The extent of cleavage of super coiled (SC) pUC 19 DNA (0.5 $\mu\text{g}$ ) to the nicked circular (NC) form was determined by agarose gel electrophoresis in Tris HCl buffer (50mM, pH 7.2) containing NaCl (50 mM). The samples were then incubated for 1 hour at 37°C followed by addition to the loading buffer containing, 25% bromophenolblue, 0.25% Xylene Cyanol, 30% glycerol and finally loaded on 0.8 % agarose gel containing 1.0  $\mu\text{g}/\text{ml}$  ethidium bromide(EB). Electrophoresis was carried out at 50V for 2 hours in Tris-borate ethylene diamine tetra acetate (TBE) buffer. Bands were visualized by UV light and photographed to determine the extent of DNA cleavage from the intensities of the bands using UVI tech Gel Documentation system. Due corrections were made for trace of NC DNA present in SC DNA sample and for the low affinity of EB binding to SC DNA in comparison to the NC form.

## RESULTS AND DISCUSSION

### Characterization of complexes

The elemental analysis, IR,  $^1\text{H}$  NMR spectral data of the new complexes are summarized in experimental section. The elemental analysis data are agreed with the theoretical values within the limit of experimental error and confirmed the formula of the complexes. These complexes are soluble in DMF, DMSO and in buffer(pH 7.2) solution. The observed conductometric measurement values in DMF solutions fall in the region 45-75  $\Omega^{-1} \text{cm}^2 \text{mole}^{-1}$  indicate their electrolytic nature. Both complexes were found to be diamagnetic.

### IR spectra

The IR spectra of ligand and complexes were recorded in the range of 4000-400  $\text{cm}^{-1}$  on KBr pellets. The IR spectra of 2-hydroxy-4-methyl-1,8-naphthyridine show bands at 1462  $\text{cm}^{-1}$  assigned to  $\nu(\text{C}=\text{N})$  aromatic hydrocarbon, 3181  $\text{cm}^{-1}$  assigned to  $\nu(\text{C}-\text{H})$  group and 3396  $\text{cm}^{-1}$  assigned to  $\nu(\text{O}-\text{H})$  group. In the spectra of complexes (1) and (2), these bands were shifted to 1636  $\text{cm}^{-1}$  and 1554  $\text{cm}^{-1}$  for  $\nu(\text{C}=\text{N})$  group, 3051  $\text{cm}^{-1}$  and 3067  $\text{cm}^{-1}$  for  $\nu(\text{C}-\text{H})$  group and 3367  $\text{cm}^{-1}$  and 3229  $\text{cm}^{-1}$  for  $\nu(\text{O}-\text{H})$  group respectively. Besides, the complexes show new bands at

410-445  $\text{cm}^{-1}$  are assigned to  $\nu(\text{M-N})$  bands [58]. In addition, the IR spectrum of the  $\text{PF}_6$  salts of each complex showed a strong band at 756  $\text{cm}^{-1}$  and 750  $\text{cm}^{-1}$  ascribed to the counter anion and this band was absent for the corresponding salts [59].

### $^1\text{H}$ NMR spectra

In the  $^1\text{H}$  NMR spectra of the complexes, the peaks due to various protons of mixed ligand of 1,10-phenanthroline and 2-hydroxy-4-methyl-1,8-naphthyridine are seen to be shifted on complexation with corresponding free ligands, suggesting complexation. The peaks due to  $\text{CH}_3$  group at 2.60 ppm, (Ar-OH) group at 8.20 ppm and (Ar-H) group at 7.80 ppm in the spectra of the ligand are shifted to 2.40 ppm, 7.80 ppm and 8.30 ppm in the spectra of complexes (1) and (2) respectively. The resonance due to (ArN-H) group at 8.80 ppm and (CH=N) group at 8.10 ppm in the spectra of the ligand are shifted to 8.70 ppm and 8.90 ppm in the spectra of complexes (1) and (2) respectively. The coordination induced shifts indicates that the ligands are coordinated to the metal ion. Based on the results above, both the complexes have octahedral geometry.





FIGURE 2. Absorption spectra of complex (1) in Tris-HCl buffer upon addition of DNA.  $[\text{Fe}] = 0.5\mu\text{M}$ ,  $[\text{DNA}] = 0\text{-}100\mu\text{M}$ . Arrow shows the absorbance changing up complex on the increase of DNA concentration. (The inset:  $[\text{DNA}]/(\epsilon_a - \epsilon_f)$  vs  $[\text{DNA}]$  for the titration of DNA with Fe(II) complex (1)).

### Absorption Spectral Studies

Electronic absorption spectroscopy is usually employed to determine the binding of complexes with DNA helix. A complex bound to DNA through intercalation is characterized by the change in absorbance (hypochromism) and red shift in wavelength, due to the intercalative mode involving a strong stacking interaction between the aromatic chromosphere and the DNA base pairs. The extent of hypochromism is commonly consistent with the strength of the intercalative

interaction [60-62].

The absorption spectra of the complexes (1) & (2) in the absence and presence of CT-DNA are given in Figures 2 and 3 respectively. Figure 2 depicts well resolved band at 322 nm for complex (1) and in Figure 3, there exist well resolved band at 328 nm for complex (2) with increasing the DNA concentration (0-100  $\mu\text{M}$ ). The result shows that the absorbance (hypochromism) decreased by the successive addition of CT- DNA to the complex solution. The hypochromism observed for the bands of complex (1) and (2) are accompanied by small bathochromic shift were 2 and 1 nm in Figures 2 and 3 respectively.



**FIGURE 3.** Absorption spectra of complex (2) in Tris-HCl buffer upon addition of DNA.  $[\text{Zn}] = 0.5\mu\text{M}$ ,  $[\text{DNA}] = 0\text{-}100\mu\text{M}$ . Arrow shows the absorbance changing up complex on the increase of DNA concentration. (The inset:  $[\text{DNA}]/(\epsilon_a - \epsilon_f)$  vs  $[\text{DNA}]$  for the titration of DNA with Zn(II) complex (2)).

The insets of figures 2 and 3 show the plots of  $[\text{DNA}]/(\epsilon_a - \epsilon_f)$  v/s  $[\text{DNA}]$  for the titration of DNA with complex (1) and complex (2) respectively. The hypochromism and bathochromic shift are observed for the complexes suggest that binding is intercalative mode. In order to compare quantitatively, the DNA binding strengths of these complexes, the intrinsic DNA binding constants  $K_b$  are determined from the decay of the absorbance at 322 nm for complex (1) and 328 nm for complex (2) with increasing concentrations of DNA. The observed  $K_b$  values for complex (1) and (2) are equal to the classical intercalators bound to CT-DNA. The  $K_b$  values for

complex (1) and (2) are  $1.8 \times 10^4 \text{ M}^{-1}$  and  $2.1 \times 10^4 \text{ M}^{-1}$  respectively. So, it is obvious that the present complexes are involved in intercalative interactions with CT-DNA.

### Viscosity Measurements

The interactions between the complex and DNA were investigated by viscosity measurements. In fact, optical photophysical probes generally provide necessary but not sufficient clues to support the binding model. Viscosity measurements that are sensitive to length change of DNA are regarded as the least ambiguous and the most critical tests of binding mode in solution in the absence of crystallographic structural data [63]. A classical intercalation mode usually resulted in lengthening the DNA helix, as base pairs were separated to accommodate the binding complex, leading to an increase in DNA viscosity. In contrast, a partial and/or non classical interaction of ligand could bend (or kink) the DNA helix, reduce its effective length and concomitantly, its viscosity [64,65]. In order to further elucidate the binding mode of the present complex, viscosity measurements were carried out on CT-DNA by varying the concentration of the added complex. As seen in Figure 4, the viscosity of DNA increased as increasing the ratio of both complexes to DNA. This result further suggested an intercalative binding mode of the complexes with DNA and also parallel to the above spectroscopic results, such as hypochromism and bathochromism of complexes in the presence of DNA.




FIGURE 4. Plot of relative viscosity Vs [complex] / [DNA]. Effect of complex (1) and complex (2) on the viscosity of CT-DNA at  $25 (\pm 0.1)^\circ\text{C}$ , [Complex] = 0-100  $\mu\text{M}$ , DNA = 50  $\mu\text{M}$ .

### Thermal Denaturation studies

The thermal behavior of DNA is a measure of the stability of DNA double helix with temperature. An interaction between DNA and complexes were indicated by the increase in the thermal melting temperature ( $T_m$ ). Thermal denaturation experiments also revealed that the intercalation of these metal complexes with DNA. In order to identify this transition process, the melting temperature  $T_m$ , which is defined as the temperature where half of the total base pairs is unbound, is usually introduced. The increase in  $T_m$  and  $\sigma_T$  of DNA could be interpreted in



terms of the stabilization that results from the intercalation of these metal complexes with DNA [66]. In the present thermal denaturation studies, thermal melting studies were carried out at DNA to complex concentration ratios of 25 and  $T_m$  and  $\sigma_T$  (the temperature range between 10% and 90% of absorption increase occurred) values were determined by monitoring the absorbance of DNA at 260 nm as a function of temperature. As shown in Figure 5, the  $T_m$  of DNA in the absence of any added drug was found to be  $65 \pm 1$  °C, under other experimental conditions. Under the same set of experimental conditions, the presence of complexes (1) and (2) increased the  $T_m$  by 4 °C and 2 °C respectively.

The observations made during the absorption titration, viscosity measurements and thermal denaturation experiments are reminiscent of those reported earlier for various metallointercalators, thus suggesting that the complexes (1) and (2) bound to DNA by intercalations [67-76].




FIGURE 5. Melting curves of CT-DNA in the absence and presence of complexes.

### DNA Cleavage studies

The oxidative DNA cleavage activity of both the complexes were studied by gel electrophoresis [77-80] using super coiled(SC) pUC19 DNA (0.5  $\mu\text{g}$ ) in Tris-HCl buffer (pH, 7.2). Figure 6 shows the gel electrophoresis separations of pUC19 DNA after 1 hour incubation with varying concentrations of the complexes. Figure 6 summarizes the results of oxidative DNA cleavage experiments carried out with the complexes (1) and (2) (at the concentration of 40  $\mu\text{M}$ , and 60  $\mu\text{M}$ ) as mentioned by the agarose gel electrophoresis method. Control experiments suggested that untreated DNA does not show any cleavage (lane 1; Figure 6). In the present study, pUC19 DNA gel electrophoresis experiment was conducted at 37 °C using the (1) and (2) in the presence of  $\text{H}_2\text{O}_2$  as an oxidant. From the results of Figure 6 at higher concentration (1) and (2) shows better nuclease activity. Control experiments using  $\text{H}_2\text{O}_2$  did not show any significant cleavage of pUC19 DNA (lane 1). At the concentration of 40  $\mu\text{M}$  and 60  $\mu\text{M}$ , complex (1) is

able to convert 75 % and 80 % of the initial SC (Form I) to NC (Form II) (lane 3). The complex (2) is able to convert 85 % (40  $\mu\text{M}$ ) and 90% (60  $\mu\text{M}$ ) of the initial SC (Form I) to NC (Form II) (lane 4 and 5).



**FIGURE 6.** Gel electrophoresis diagram showing the cleavage of SC pUC 19 DNA (0.5  $\mu\text{g}$ ), by the complexes (1) and (2) in a buffer containing 50 mM Tris-HCl and 50 mM NaCl 37 °C in the presence of  $\text{H}_2\text{O}_2$ : lane 1. DNA control; lane 2. DNA+40  $\mu\text{M}$  (1); Lane 3. DNA+60  $\mu\text{M}$  (1); Lane 4. DNA+40  $\mu\text{M}$  (2); Lane 5. DNA+60  $\mu\text{M}$  (2).

## CONCLUSION

In conclusion, we have synthesized and characterized two new complexes of the type  $[\text{Fe}(\text{L}_2)_2\text{L}_1](\text{PF}_6)_2$  [Complex (1)] and  $[\text{Zn}(\text{L}_2)_2\text{L}_1](\text{PF}_6)_2$  [Complex (2)]. Interactions of the new complexes with (double stranded) DNA were investigated by absorption spectra, viscosity and thermal denaturation studies. From the experimental results, it was confirmed that the complexes bound with the double stranded DNA with binding constant  $K_b = 1.8 \times 10^4 \text{ M}^{-1}$  for Complex (1) and  $K_b = 2.1 \times 10^4 \text{ M}^{-1}$  for Complex (2), respectively. The viscosity of solution of the DNA bound to the complexes increased with increase in concentration of the complexes. Thermal denaturation experiments revealed the intercalation of both complexes with DNA. Further, the cleavage studies show that the complex (1) and (2) have significant nuclease activity.

## Acknowledgement

The authors are thankful to Sri.B.Premanath Reddy, Chairman, Acharya Institutes and Prof. G.P.Prabhukumar, Director, Acharya Institute of Technology, Bangalore, India.

## REFERENCES

- [1] N. A. P. Kane-Maguire and J. F Wheeler, *Coord Chem Rev.*, **2001**, 211, 145.
- [2] C. Kaes, A. Katz, and M. W. Hosseini, *Chem Rev.*, **2000**, 100, 3553.
- [3] B. Armitage, *Chem Rev.*, **1998**, 98, 1171.
- [4] D. S. Sigman, A. Mazumder, and D. M. Perrin, *Chem Rev.*, **1993**, 93, 2295.
- [5] M. K. De-Armond and C. M. Carlin, *Coord Chem Rev.*, **1981**, 36, 325.
- [6] B. H. Geierstanger, M. Marksich, P. B. Dervan, and D. E. Wemmer, *Science.*, **1994**, 266, 646.
- [7] C. Liu, J. Li-Zhou, L. Liao-Wang, and Z. H. Xu, *J Inorg Biochem.*, **1996**, 75, 233.
- [8] G. Prativiel, J. Bernadou, and B. Meunier, *Angew Chem Int Ed Engl.*, **1995**, 34, 746.
- [9] S. O. Kelly and J. K. Barton, *Metal ions in biological systems* (Eds. A. Siegel and H. Sigel) Marcel Dekker, New York, **1999**, 39, 211.

- [10] B. Norden, P. Lincoln, B. Akerman and E. Tuite, *Metal ions in biological systems* (Eds. A. Siegel and H. Sigel) Marcel Dekker, New York, **1996**, 33, 177.
- [11] A. K. Mesmaeker, J. P. Lecomte, and J. M. Kelly, *Topics in Curr Chem.*, **1996**, 177, 25.
- [12] D. S. Sigman, A. Mazumder, and D. M. Perrin, *Chem Rev.*, **1993**, 93, 2295.
- [13] R. J. Morgan, S. Chatterjee, A. D. Baker, and T. C. Streckas, *Inorg Chem.*, **1991**, 30, 2687.
- [14] S. T. Frey, H. H. J. Sun, N. N. Murthy, and K. D. Karlin, *Inorg Chim Acta.*, **1996**, 242, 329.
- [15] A. S. Sitlani, E. C. Long, A. M. Pyle, and J. K. Barton, *J Am Chem Soc.*, **1992**, 114, 2303.
- [16] J. K. Barton, *Science*, **1986**, 233, 727.
- [17] M. Carter, M. Rodriguez, and A. J. Bard, *J Am Chem Soc.*, **1989**, 111, 8901.
- [18] R. Vijayalakshmi, M. Kanthimathi, V. Subramanian, and B. Nair, *Biochem Biophys Res. Commun.*, **2000**, 271, 731.
- [19] V. V. Ganesan, and B. U. Nair, *J Inorg Biochem.*, **2003**, 93, 271.
- [20] C. Turro, S. H. Bossman, Y. Jenkins, J. K. Barton, and N. J. Turro, *J Am Chem Soc.*, **1995**, 117, 9026.
- [21] X. H. Zou, B. H. Ye, H. Li, J. G. Liu, Y. Xiong and L. N. Ji, *J Chem Soc, Dalton Trans.*, **1999**, 1423.
- [22] Y. Xiong and L. N. Ji, *Chem Rev*, **1999**, 185, 711.
- [23] Y. Xiong, X. F. He, X. H. Zou, J. Z. Wu, X. M. Chen, L. N. Ji, R. H. Li, J. Y. Zhou and K. B. Yu, *J Chem. Soc. Dalton Trans.*, **1999**, 19.
- [24] J. G. Liu, B. H. Ye, Q. L. Zhang, X. H. Zhou, X. Tian and L. N. Ji, *J Biol Inorg Chem.*, **2000**, 5, 119.
- [25] J.P. Rehmman, and J. K. Barton, *Biochemistry*, **1990**, 29, 1701.
- [26] S. Arounaguirri and B. G. Maiya, *Inorg Chem.*, **1996**, 35, 4267.
- [27] L. Jin and P. Yang, *Polyhedron*, **1997**, 16, 3395.
- [28] X. F. He, L. Wang, H. Chen, L. Xu and L. N. Ji, *Polyhedron*, **1998**, 17, 3161.
- [29] Q. L. Zhang, J. G. Liu, H. Chao, G. Q. Xue and L. N. Ji, *J Inorg Biochem.*, **2001**, 83, 49.
- [30] S. Arounaguirri and B. G. Maiya, *Inorg Chem.*, **1999**, 38, 842.
- [31] C. J. Burrows and S. E. Rokita, *Acc Chem Res.*, **1994**, 27, 295.
- [32] A. M. Pyle, J. K. Barton and S.J. Lippard, Ed. *Progress in Inorganic Chemistry*. Wiley, New York, **1990**, 38, 413.
- [33] T. D. Tullius, *Metal-DNA Chemistry*. ACS Symposium Series, *American Chemical Society*, Washington, DC, Vol. 402, **1989**.
- [34] A. R. Banerjee, J. A. Jaeger and D. H. Turner, *Biochemistry*, **1993**, 32, 153.
- [35] S. Ramakrishnan and M. Palaniandavar, *J Chem Sci*, **2005**, 117, 179.
- [36] P. Lu, M. L. Zhu and P. Yang, *J Inorg Biochem*, **2003**, 95, 31.
- [37] C. Hemmert, M. Pitie, M. Renz, H. Gornitzka, and S. B. Meunier, *J Biol Inorg Chem.*, **2001**, 6, 14.
- [38] M. Navarro, E. J. Cisneros-Fajardo, A. Sierralta, M. Fernandez-Mastre, P. Silva, D. Arrieche and E. Marchan, *J Biol Inorg Chem.*, **2003**, 8, 401.
- [39] H. J. Epply, S. M. Lato, A. D. Ellington, and J. M. Zaleski, *Chem Commun.*, **1999**, 2405.
- [40] S. Dhar, D. Senapati, P. K. Das, P. Chattopadhyay, M. Nethaji, and A.R. Chakravarthy, *J Am Chem Soc.*, **2003**, 125, 12118.
- [41] P. J. Benites, R. C. Holmberg, D. S. Rawat, H. H. Thorp, and J. M. Zaleski, *J Am Chem Soc.*, **2003**, 125, 6434.
- [42] T. M. Bray and W. J. Bettger, *Biol Med.*, **1990**, 8, 281.
- [43] S. R. Powell, *J Nutr.*, **2000**, 130, 1447.

- [44] I. E. Dreosti, *Mutate Res.*, **2001**, 475, 161.
- [45] R. J. Cousins, *Proc Nutr Soc.*, **1998**, 57, 307.
- [46] K. H. Falchuk, *Mol Cell Biochem.*, **1998**, 188, 41.
- [47] A. S. Prasad, *J Am. Coll Nutr.*, **1998**, 17, 542.
- [48] A. S. Prasad, and O. Kucuk, *Cancer Metastasis Rev.*, **2002**, 21, 291.
- [49] A. S. Prasad, *BMJ.*, **2003**, 326, 409.
- [50] D. S. Sigman, A. Mazumder and D. M. Perrin, *Chem Rev.*, **1993**, 96, 2295.
- [51] T. Ghosh, B. G. Maiya and A. Samanta, *J Biol Inorg Chem.*, 2005 10, 496.
- [52] N. J. Turro, J. K. Barton and D. A. Tomalia, *Acc Chem Res.*, **1991**, 24, 332.
- [53] F. Q. Liu, Q. X. Wang, K. Jiao, F. F. Jian, G. Y. Liu, R. X. Li, *Inorg. Chim. Acta.*, **2006**, 395, 1524.
- [54] T. F. Tullis, *Metal-DNA chemistry*, ACS Symposium Series, Vol. No. 402, American Chemical Society, Washington DC, USA, **1989**.
- [55] J. K. Barton, *Science*, **1986**, 233, 727.
- [56] M. E. Reichmann, S. A. Rice, C. A. Thomas and P. Doty, *J Am. Chem. Soc.*, **1954**, 76, 3047.
- [57] A. Wolfe, G. H. Shimer and T. Meehan, *Biochemistry*, **1987**, 26, 6392.
- [58] R. Augamuthu, V. Rajendiran, P. UmaMaheshwari, R. Balamurugan, C. A. Kilner, M. A. Halcrow and M. Palaniandavar, *J Inorg Biochem.*, **2005**, 99, 1717.
- [59] H. Zhang, C. S. Liu, X. H. Bu and M. Yang, *Journal of Inorg Biochem.*, **2005**, 99 1119.
- [60] C. N. Sudhamani, H. S. Bhojya Naik, T. R. Ravikumar Naik and M. C. Prabhakar, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, **2009**, 72, 643.
- [61] L. Mishra, K. Y. Ajay, S. P. Ratna C. S. Choi and A. Koji, *Metal Based Drugs*, **2001**, 8, 65.
- [62] J. K. Barton, A. T. Danishefsky and J. M. Goldberg, *J Am. Chem. Soc.*, **1984**, 106, 2172.
- [63] S. A. Tysoe, R. J. Morgan, A. D. Baker and T. C. Streckas, *J Phys. Chem.*, **1993**, 97, 1707.
- [64] L. Jie, T. B. Lu, H. Li, Q. L. Zhang, L. N. Ji, T. X. Zhang, L. H. Qu and H. Zhou, *Transition Metal Chemistry*, **2002**, 27, 686.
- [65] K. Laxma Reddy, Y. Harish Kumar Reddy and S. Satyanarayana, *Nucleosides, Nucleotides and Nucleic Acids*, **2009**, 28, 953.
- [66] H. Chao, Y. X. Yuan, F. Zhou, L. N. Ji and J. Zhang, *Transition Metal Chemistry*, **2006**, 31, 465.
- [67] C. B. Archika, G. Smita, A. K. Anupa, S. K. Avinash, and G. Vedavati, *Transition Metal Chemistry*, **2005**, 30, 312.
- [68] M. Surendrababu, P. G. Krishna, K. Hussain Reddy and G. H. Phillip, *Main Group Chemistry*, **2009**, 103, 8.
- [69] S. Anbu, M. Kandaswamy and B. Varghese, *Dalton Trans.*, **2010**, 39, 3823.
- [70] B. Sreekanth, G. Krishnamurthy, H. S. Bhojya Naik, M. C. Prabhakara, T. K. Vishnuvardhan, *Synthesis and Reactivity in Inorganic, Metal-Organic and Nano-Metal chemistry*, **2010**, 40, 955.
- [71] C. B. Spillane, M. N. V. Dabo and N. C. Fletcher, *J Biol. Inorg. Chem.*, **2008**, 102, 673.
- [72] M. Surendrababu, P. G. Krishna, K. Hussain Reddy and G. H. Phillip, *J Serb Chem Soc.*, **2010**, 75, 61.
- [73] B. Peng, C. Xiang, K. J. Du, B. L. Yu, H. Chao and L. N. Ji, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, **2009**, 74, 896.
- [74] A. Arounaguirri, D. Eshwaramoorthy, A. Ashokkumar, A. Dattagupta and B. G. Maiya, *Proceedings of the Indian Academy of Sciences: Chemical Sciences*, **2000**, 112, 1.

- [75] P. Nagababu, J. Naveena Lavanya Latha, Y. Prashanthi, S. Satyanarayana, *Journal of Chemical and Pharmaceutical Research*, **2009**, 1, 238.
- [76] Penumaka Nagababu, J. Naveena Lavanya Latha, Mynam Shilpa, P. Pallavi, Rajender Reddy, C. Shobha Devi, S. Satyanarayana, *Journal of Chemical and Pharmaceutical Research*, **2010**, 2, 144.
- [77] M. C. Prabhakar, H. S. Bhojya Naik, *Biometals*, **2008**, 21, 675.
- [78] B. Sreekanth, G. Krishnamurthy, H. S. Bhojya Naik, T. K. Vishnuvardhan, *Nucleosides, Nucleotides and Nucleic Acids*, (**2011**) (In Press).
- [79] M. C. Prabhakar, H. S. Bhojya Naik, V. Krishna, H. M. Kumaraswamy, *Nucleosides, Nucleotides, and Nucleic Acids*, **2007**, 26, 459.
- [80] B. Sreekanth, G. Krishnamurthy, H. S. Bhojya Naik, T. K. Vishnuvardhan, B. Vinaykumar, N. Sharath, *Nucleosides, Nucleotides and Nucleic Acids*, **2011**, 30, 83.