



Research Article

ISSN : 0975-7384
CODEN(USA) : JCPRC5

GC-MS analysis of bioactive compounds in phenolic extracts of leaves and flowers of *Chrysanthemum cinerariaefolium* and their efficacy against larvae of *Culexquin quefaciatus* Say(Diptera: Culicidae)

Ruqia O. Kareem and Mohammed R. Annon

Biology Department, College of Science, AL- Qadisiyah University, AL- Qadisiyah, Diwania, Iraq

ABSTRACT

The bioactive compounds of phenolic extract of leaves and flowers of (pyrethrum) *Chrysanthemum cinerariaefolium* have been investigated and evaluated using GC-MS analysis. (Perkin Elmer GC clarus 500) Gas Chromatography - Mass Spectrometry, The mass spectra of bioactive compounds were matched according to National Institute of Standards and Technology (NIST) library. The phenolic extract of leaves contains four compounds, while the flowers extract has 13 compounds, The analysis showed also that 3,7,11,15-Tetramethyl-2-hexadecen-1-ol, 3-Buten-2-one, and 4-(2-hydroxy-2,6,6-trimethylcyclohexyl) were found in both leaves and flowers extract. Then bioassay of these two phenolic extract were tested against all larval instar of *Cx. quinquefaciatus* and showed that the first larval instar was more sensitive than other preceding instars and the mortality percentages of these larval instars were high in phenolic extract of flowers.

Keywords: GC-MS, *C. cinerariaefolium*, Phenolic compounds, *Cx. quinquefaciatus*

INTRODUCTION

Culex quinquefaciatus is distributed abundantly in Iraq, especially in the central and southern parts, This species is vector of some pathogens such as *Wuchereriabancrofti* ,viruses such as causing West Nile Fever, Japanese Encephalitis as well as Dengue Fever [1],[2]. Botanical pesticides is one of promising caudate to control vectors disease[3]and are environmentally safe, degradable and target-specific[4]. *C.cinerariaefolium* It is an important source of pyrethrins [5],[6]. Phenols are harmful because of their ability to join with proteins by hydrogen bonds.[7]

EXPERIMENTAL SECTION

Insect cultuer

The cultuer of *Culex quinquefaciatus* were maintained at entomology laboratory, college of science, Qadisiyah University, at a temperature of 28C ° with a relative humidity of 75C ° ± 5 % for 14-h photoperiod. The larvae of *Cx. quinquefaciatus* were maintained in separate enamel containers.

Plant samples

Leaves and flowers of the plant were collected in the early morning (before sunrise), and then were washed and cleaned and dried naturally at lab temperature. The dry leaves and flowers were ground separately using an electric grinder and kept in dark plastic containers ready for use.

Preparation of crude phenolic extract

The phenolic extract was attended according to [8] as follows: 20 gms of dried powder of leaves and flowers of *C.cinerariaefolium* was blended separately 400 ml solution of concentrated hydrochloric acid HCL 36% (2ml).

Extraction was done using reflex condenser in a water bath of boiling water (100 °C) for one hour, then left until cool and then filtered through Whatman No.1 filter paper. The filtrate was transferred to the separating funnel and the same volume of n- propanol was added. Then a sufficient quantity of NaCl was added until saturation. After separation the top layer (phenolic) taken and dried in rotary evaporator at 45C°, and then left to dry at lab. temperature. The resulting product was then collected and kept in sealed glass tubes in the refrigerator at 4 C° ready for use.

GC-MS method

GC-MS analysis was carried out on a GC clarus 500 PerkinElmer system comprising a AOC-20i auto sampler and gas chromatograph interfaced to a mass spectrometer instrument under the following conditions: column Elite-1 fused silica capillary column (30 x 0.25 mm ID x 1 μ Mdf, composed of 100% Dimethyl polysiloxane), operating in electron impact mode at 70eV; Helium gas (99.999%) was used as carrier gas at a constant flow of 1 ml /min and an injection volume of 0.5 μI was employed (split ratio of 10:1) injector temperature 250 °C; ion-source temperature 280 °C. The oven temperature was set at 110 °C (isothermal for 2 min), with an increase of 10 °C/min, to 200°C, then 5 °C/min to 280 °C, ending with a 9min isothermal at 280 °C. Mass spectra were taken at 70eV; a scan interval of 0.5 seconds and fragments from 40 to 450 Da. Total GC running time was 36min. The relative percentage amount of each component was calculated by comparing its average peak area to the total areas. The software adapted to handle mass spectra and chromatograms was Turbo Mass Ver 5.2.0 [9].

Identification of components

Interpretation of mass spectrum GC-MS was conducted using the database of National Institute Standard and Technology(NIST) having more than 62,000 patterns. The spectrum of the unknown component was compared with the spectrum of the known components stored in the NIST library. The name, molecular weight and structure of the components were ascertained.

Bioassay of phenolic extract

In order to estimate the biological activity of the extract of crude phenolic compounds of *C.cinerariaefolium* leaves and flowers was attended the Stock solution by dissolving 4 gms of dry matter for each of the leaves, flowers separately in 20 ml of ethanol 96% and finished size to 100 with distilled water, bringing to the Stock solution 4% or (40 mg /ml), and from the last solution attended concentrations (2.5, 5, 10, 20, 40) . The control treatment was 5ml ethanol added to it 95 ml of distilled water. The experimental tubes was 3 replications each containing 100 ml from each concentration and explained where 20 larvae were mortality ratio calculated after 24 hours. and corrected ratios according to the equation and subjected all experiments for statistical analysis by using Least significant Differences (L.S.D) $p \leq 0.05$ using Abbott's formula [1925][10].

$$\% \text{ mortality} = \frac{\% \text{ mortality in treatment} - \% \text{ mortality in control}}{100 - \% \text{ mortality in control}} \times 100$$

RESULTS AND DISCUSSION

Four phenolic compounds were identified in *C. cinerariaefolium* leaves and thirteen compounds were also identified in flowers. The name, retention time (RT), molecular formula, molecular weight(MW) are presented in (Table1, and Fig.1, Table 2, Fig.2). Also it turns out that some of the compounds were present in all of the leaves and flowers, but differ in the retention time, such as 3,7,11,15-Tetramethyl-2-hexadecen-1-ol, 3-Buten-2-one, 4-(2-hydroxy-2,6,6-trimethylcyclohexyl)-.

Table 1: GC-MS analysis of *Chrysanthemum cinerariaefolium* leaves extract

Peak #	R. time	Area %	Name	Molecular formula	Molecular weight
1	12.62	10.28	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C ₂₀ H ₄₀ O	296
2	13.56	7.72	1,1'-Bicyclohexyl, 2-(2-methylpropyl)-, trans-	C ₁₆ H ₃₀	222
3	18.57	76.07	3-Buten-2-one, 4-(2-hydroxy-2,6,6-trimethylcyclohexyl)-	C ₁₃ H ₂₂ O ₂	210
4	20.42	5.92	1,2-Benzenedicarboxylic acid, diisooctyl ester	C ₂₄ H ₃₈ O ₄	390




Fig.5: Chromatogram obtained from the GC/MS leaves extract of *C. cinerariaefolium*

Chemical detection using GC-MS demonstrated that the plant extract of *Boerhavia diffusa L.* contains Steroles, Tannins and Flavonoids as well as Phenolic compounds [11]. [12] selected methanol extract compounds of roots, stems and leaves of *Labisia paucifolia*, where the results indicate that the leaves extract contains a large amount of phenolic compounds and flavonoids compared with the roots and stems extract, and was all of gallic acid, Kaempferol are more flavonoids and phenols presence.

Table 2: GC-MS analysis of *Chrysanthemum cinerariaefolium* flowers extract

Peak #	R. time	Area %	Name	Molecular formula	Molecular weight
1	5.87	28.07	Acetic acid	C ₂ H ₄ O ₂	60
2	6.98	7.11	Camphor	C ₁₀ H ₁₆ O	152
3	9.39	8.04	Bicyclo[2.2.1]heptane, 2-methoxy-1,7,7-trimethyl-	C ₁₁ H ₂₀ O	168
4	10.45	2.01	2-Butenoic acid, 2-methyl-, (Z)-	C ₅ H ₈ O ₂	100
5	11.63	1.77	1-Hexadecanol	C ₁₆ H ₃₄ O	242
6	12.58	7.22	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C ₂₀ H ₄₀ O	296
7	13.31	1.39	1,3-Cyclopentadiene, 5,5-dimethyl-2-ethyl-	C ₉ H ₁₄	122
8	13.60	6.10	Diazoacetic acid, 2-isopropyl-5-methylcyclohexyl ester	C ₁₂ H ₂₀ N ₂ O ₂	224
9	16.87	5.68	3,6-Octadien-1-ol, 3,7-dimethyl-, (Z)-	C ₁₀ H ₁₈ O	154
10	17.22	2.93	3-Oxabicyclo[4.1.0]heptan-2-one, 4,4,7,7-tetramethyl-	C ₁₀ H ₁₆ O ₂	168
11	18.00	2.20	12-Oxabicyclo[9.1.0]dodeca-3,7-diene, 1,5,5,8-tetramethyl-, [1R-1R*,3E,7E,11R*]-	C ₁₅ H ₂₄ O	220
12	18.42	14.81	3-Buten-2-one, 4-(2-hydroxy-2,6,6-trimethylcyclohexyl)-1	C ₁₃ H ₂₂ O ₂	210
13	22.88	12.66	Thujone	C ₁₀ H ₁₆ O	152




Fig.6: Chromatogram obtained from the GC/MS with the extract of *C. cinerariaefolium* flowers

GC-MS analysis used to explained that content of plant *Dolichandrone atrovirens* leaves extract , where it found container on saponins, phenols, flavonoids and vitamin C, so that this plant use of pharmaceuticals and pharmaceuticals purposes [13]. Phenols are the secondary metabolites that a ubiquitously present in plants. They have been suggested to play a role in the antioxidant function . Phenolic compounds have antioxidant properties because of their ability to scavenge free radicals.

The phenolic compounds of plant origin showed their antioxidative effect by various mechanisms, including their ability to scavenge free radicals or activate various antioxidant enzymes and inhibit oxidizes[14].[15]investigated the presence of phytochemical contents of the selected medicinal plants. Proteins, carbohydrates, phenols, tannins, flavonoids, saponins were detected in all of the plants tested.[16] was determined the larvicidal activity of the ethanol extract of *E. rutaecarpa* unripe fruits and the isolated constituents against the larvae of the Culicidae mosquito *Ae. albopictus*. The presence of phytochemical compounds of carbohydrates, terpenoids, steroids, flavonoids, phenols was screened by qualitative method.[17] reported 49 phytoconstituent in ethanolic extract of *Maranta arundinacea*. L subjected to GC MS analysis.

BIOASSAY

The phenolic extract of flowers record high mortality was 77.79 % .The results also indicate the sensitivity of the larval instar to ward the extract . Where the first instar was the most sensitivity to the extract and in the all concentration used compared to the other larval stages .The first stages record ratios of mortality of (77.33-84.24)% compared tp the second stages (68 .10 – 80. 54)% , third (64. 66 – 74.22)% and fourth (62.99 – 72.47) in the phenolic extract of leaves and flowers respectively.




Fig. 1: effects of phenolic compounds extract of the leaves and flowers of *C. cinerariaefolium* on mortality of first larval instar of *Cx. quinquefaciatus*




Fig. 2: effects of phenolic compounds extract of the leaves and flowers of *C. cinerariaefolium* on mortality of second larval instar of *Cx. quinquefaciatus*




Fig. 3: effects of phenolic compounds extract of the leaves and flowers of *C. cinerariaefolium* on mortality of third larval instar of *Cx. quinquefasciatus*




Fig. 4: effects of phenolic compounds extract of the leaves and flowers of *C. cinerariaefolium* on mortality of fourth larval instar of *Cx. quinquefasciatus*

Phenolic compounds cause two types of physiological effects in the larvae tissues were toxic effect in directly disrupt secretion in the nervous system, or a direct effect through the spread of the entry into force of these compounds in the target tissue [18],[19]. It may return the cause of the effectiveness of phenols on other secondary metabolites to the different active substances in the *C. cinerariaefolium* plant and it contains tannins that are toxic compounds to insects as associated with saliva and digestive enzymes including trypsin and chymotrypsin and then inhibited and thus the insects start to lose weight and then death [20].

Acknowledgements

The authors are gratefully thank to Dr. Salwam Ali, ecology Department and Haider R. Saud, chemistry Department, college of Science, Al-Qadisiyah university for their helps to complete this work successfully.

REFERENCES

- [1] KSTennyson; J Ravindran and SARivoli. *Asi. Paci. J. Trop. Bio. Med.*, **2012**, 2(2): S1130-S1134.
- [2] PKareru; ZK Rotich and ELMaina. *The Afri. Case, Insecticides - Development of Safer and More Effective Technologies*. **2013**. 299 PP.
- [3] FYarahmadi; ARajabpour; NZSohani and LRamezani. *Avi. J. of Phytomed.* **2013**, 3(2): 106-111.
- [4] S Senthil Nathan; and K Saehoon. *Crop Prot.*, **2006**, 25(3): 287-291.
- [5] AHitmi; ACoudret and CBarthomeuf. *Crit. Rev. Biochem. Mol. Biol.* **2000**, (35): 317-337.
- [6] CBisht; A Badoni; RKVashishtha and MCNautiyal. **2009**. 5(4): 147 -150.

- [7] RS Rattan. Mechanism of action of insecticidal secondary metabolites of plant origin. (2010). Crop Prot. 29 : 913-920.
- [8] JB Harborne. phytochemical methods. Chapman and Hall. New York 2nd Ed. 1984. 288pp.
- [9] K Srinivasan; SSivasubramanian and SKumaravel. *Int. J. Pharm. BioSci.* **2013**, 5(1):714-720.
- [10] W SAAbbott. *J. Econ. Entomol.* **1925**, 18: 265-267.
- [11] GRJ Beegum; SSBeevy and SVSugunan. *Int. J. of Emer. Techn. and Advan. Eng.* **2014**, (7):318-324.
- [12] HZEJaafar; EKarimi; MH Ibrahim and AGhasemzadeh. *aust. J. of crop Sci.* **2014**, 7(2):276-280.
- [13] SKayarohanam and SKavimani. *Int. J. Pharm. Res.* **2015**, 6(3):219-222.
- [14] WLeslie and KBoxin. *J. Agric. Food. Chem.* **2002**, 50:3495-3500.
- [15] RYadav and M Agarwala. *J. Phyto.* **2011**, 3(12):10-14.
- [16] Z LLiu; Q Z Liu; S SDuan and Z W Deng. *Paras. Res.* **2012**, 111(3):991-996.
- [17] SNishaa; MVishnupriya; JM Sasikumar and VKGopalakrishnan. *L. Res. J. Pharma, Biol. Chem. Sci.*, **2013**, 4(2):52-59.
- [18] RFChapman. The insect structure and function. *The Eng. Univ-press*, **1978**. 670pp.
- [19] B C Freeman; and G A Beattie. An overview of Plant Defenses against Pathogens and Herbivores The Plant Health Instructor. DOI: 10.1094 /PHII-1- . **2008**. 226.
- [20] RKSingh; R CDhiman. and PKMittal. *J. Vect. Born. Dis.*, **2006**. 43(2) 88-91.