



Research Article

ISSN : 0975-7384
CODEN(USA) : JCPRC5

Synthesis, antibacterial and antifungal properties of newer (2-butyl-5-amino-3-benzofuranyl)-4-methoxy phenyl methanone heterocyclic and isoxazolic acids

Yogesh D. B.^{1*}, Dinesh B. Shenoy¹, Sunil Kumar Agarwal¹, Menaka T.³, Srinivasa G. R.², Nagaraja G.¹ and Sunil Kumar U. S.³

¹Department of Pharmaceutical Chemistry, Priyadarsini College of Pharmacy and Research Centre, under NIMS University, (Rajasthan, Jaipur), Hanumanthapura, Koratagere, Tumkur (D), Karnataka (S), India

²Alkem Laboratories, Peenya Industrial Area, Bangalore, Karnataka(S), India

³Department of pharmaceutical Analysis, Mallige College of Pharmacy, Siluvepura, Bangalore, Karnataka(S), India

ABSTRACT

The (2-butyl -5-amino-3-benzofuranyl) 4-methoxy phenyl methanone heterocyclic and isoxazolic acids have been synthesized by the reaction of (2-butyl -5-amino-3-benzofuranyl) 4-methoxy phenyl methanone with different heterocyclic and isoxazolic acids were described (DBMY20-28). All the compounds were synthesized by conventional method and characterized by IR, ¹H NMR and mass spectral data. The synthesized compounds were tested for antimicrobial activity against *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa*. Among the synthesized compounds (2-butyl -5-amino-3-benzofuranyl) 4-methoxy phenyl methanone acids (DBMY-22,27,28) was found the most active derivative against *S. aureus*, *E. coli* and *P. aeruginosa* and The compounds DBMY-22, 27, 28 was found the most active derivative against *Candida albicans* and *Asperigillus fumigatus* respectively. The other Compounds exhibited moderate activity against the other test microorganisms.

Keywords: Antimicrobial activity, conventional, Amino Benzofuran, Heterocyclic and Isoxazolic acids.

INTRODUCTION

It is an object of this invention to provide new medicinal compounds. This invention relates to new Benzofuran derivatives, their preparation, characterization and biological use. Upon further study of the specification and appended claims, further objects and advantages of this invention will become those skilled in the art. Benzofuran derivatives exhibit various types of physiological activity and enter into medicinal products and this determines the great attention that has been paid to the synthesis of new compounds of this series [1-2].

The Benzofuran derivatives drew a lot of attention on various pharmacological activities like anti-microbial[3-5], anti-inflammatory[6], antioxidant[7], antithyroid[8] and antitumour[9-11] and in the treatment of allergic conditions, asthma, cardiovascular disorders and as cytoprotective agents[12], arrhythmia[13] and Alzheimer diseases[14-15]. Benzofurans have been drawn as promising structural units in the field of medicinal chemistry. Ameridone, Dronedarone, are well known marketed drugs. Owing to the importance and established pharmacological activity of these compounds we directed our attention towards synthesis and study of biological activity of some newer derivatives of Benzofuranamine substituted compounds.

EXPERIMENTAL SECTION

Thin layer chromatography was used to monitor the completion of the reaction and homogeneity of the synthesised compound. Melting points were determined using a manual Buchi electro thermal apparatus (range 0-300 c) in open capillary tubes and uncorrected. IR spectra in KBr pellets were recorded Perkin-Elmer Spectrum 100 FT IR spectrometer (400 MHz) in DMSO-d₆ /CDCl₃ using TMS as an internal standard (chemical shifts are expressed in ppm). The homogeneity of the compounds was checked on silica gel-G coated plates, Hexane, ethyl acetate and Chloroform as the eluent and observed in UV lamp, iodine vapours or KMnO₄ spray as developing agents. All the synthesised compounds gave satisfactory elemental analysis.

Synthesis of 2-hydroxy-5-nitro phenylphosphonium bromide. [16]

100 g(0.43 mole) of 2-hydroxy 5-nitrobenzyl bromide and 113 g (0.43 mole) of triphenylphosphine are heated at reflux for 0.5 hour in 1600 ml of chloroform. The mixture is allowed to cool and the white precipitate formed is filtered off. The filtrate is evaporated to dryness in a vacuum and the residue is taken up in 500 ml of toluene. The precipitate is filtered off, washed with toluene and the solids formed are pooled and dried in a vacuum at 50 C. In this manner, 210.5 g of 2-hydroxy 5-nitrobenzyl triphenylphosphonium bromide are obtained. Yield: 99.01%

Synthesis of 2-butyl 5-nitro Benzofuran.


48.98 g (0.94 mole) of pentanoyl chloride are added slowly with stirring to a mixture of 200g (0.40 mole) of 2-hydroxy 5-nitrobenzyltriphenylphosphonium, bromide and 120.2g (1.52 mole) of pyridine in 700 ml of chloroform. The mixture is heated at reflux for 2 hours. 2800 ml of toluene is added and 1400 ml of solvents are distilled. 228 g (2.28 moles) of triethylamine are then added and the mixture is heated at reflux for 3 hours. It is allowed to cool, the triphenylphosphine-oxide, formed is filtered off, washed with ethyl acetate and the filtrate is concentrated in a vacuum. The viscous residue obtained is dissolved in acetonitrile and extracted with pentane in a liquid-liquid, extraction apparatus. The solution is stirred over sodium sulphate, filtered and evaporated to dryness. In this manner, crude 2-n-butyl 5- nitrobenzofuran is obtained. Purity [high performance liquid chromatography] (HPLC):97.9% B.p.120-123 C.

Synthesis of (2-butyl -5-nitro-3-benzofuranyl) 4-methoxy phenyl methanone.

Tin chloride (2.7ml, 23.0mmol) was added drop wise to a stirred solution of N-2-butyl 5-nitrobenzofuran (2.50g, 9.4mmol) and P-anisoyl chloride (3.62g, 10.0mmol) in dichloromethane (20ml) at 0-5 C. The mixture was stirred for 30 min at the same temperature and for 2 hours at room temperature. Water (50ml) was added drop wise to the stirred mixture at 0-5 C. the mixture was extracted with dichloromethane (3X30 ml). the combined organic layers were washed with saturated sodium bicarbonate aqueous solution, dried over sodium sulphate and evaporated .3.45g (65.9%) of (2-butyl -5-nitro-3-benzofuranyl) 4-methoxy phenyl methanone was obtained after purification of the residue by column chromatography on silica gel 60-120(ethyl acetate/hexane 1:3v/v). LC-MS: m/z 354.15 [M+]. IR (KBR): 1720,1545,1465,1370 Cm⁻¹, ¹H NMR(CDCl₃): δ 0.92-1.01 (3H,t,-CH₃-); δ 1.38-1.40 (2H,m-CH₂-); δ 1.58-1.64 (2H,m,-CH₂-); δ 1.99-2.22 (2H,t,-CH₂-); δ 3.82-3.90 (3H,s,-Ar-OCH₃-); δ 7.02-8.56 (7H,m,-Ar-).Anal. Calcd for C₂₀H₁₉NO₅: C, 67.98; H, 5.42; N, 3.96. Found: C, 66.90; H, 5.62; N, 3.68.


Synthesis of (2-butyl -5-amino-3-benzofuranyl) 4-methoxy phenyl methanone (5A).

Dissolved (2-butyl -5-nitro-3-benzofuranyl) 4-methoxy phenyl methanone in 10 ml of methanol added sodium formate and zinc dust .stirred for 2-3 hour until greenish yellow colour. The progress of the reaction was monitored by the TLC. Reaction mixture was distilled out under reduced pressure. The (2-butyl -5-amino-3-benzofuranyl) 4-methoxy phenyl methanone was purified by column chromatography using silica gel 60-120(hexane/ethyl acetate7:3) to remove starting material, followed by (chloroform /methanol 9:1).yield is 90% purity: 98.70%. LC-MS: m/z 324.25 [M+]. IR (KBR): 3450,1720,1465,1370,1290 Cm⁻¹, ¹H NMR(CDCl₃): δ 1.00-1.20 (3H,t,-CH₃-); δ 1.42-1.48 (2H,m-CH₂-); δ 1.80-1.86 (2H,m,-CH₂-); δ 2.30-2.46 (2H,t,-CH₂-); δ 3.79-3.92 (3H,s,-Ar-OCH₃-); δ 4.22-4.38 (3H,s,-Ar-NH₂-); δ 6.932-8.77 (7H,m,-Ar-).Anal. Calcd for C₂₀H₂₁NO₃: C, 74.28; H, 6.55; N, 4.33. Found: C, 74.00; H, 6.72; N, 4.82.




Scheme-1. [16]

Scheme-2

STEP 1: Formation of acid chloride

Heterocyclic/isoxazolic carboxylic acid was dissolved in dichloromethane added 2-3 ml of dimethyl formamide and one equivalent of oxalyl chloride at 5-10°C stirred for 30 minutes and reaction was monitored by TLC and reaction mixture taken for next stage.

STEP 2: Acid Chloride to Amide.

Distilled out the solvent from the step one to remove excess of oxalyl chloride. Acid chloride (1mmol) and (2-butyl - 5-amino-3-benzofuran-3-yl) 4-methoxy phenyl methanone (1mmol) was dissolved in dichloromethane (10ml), triethylamine (1mmol) stirred for 2 hours and reaction was monitored by TLC and purified by column chromatography using silica gel 60-120. yield 80%. purity -99%.

Synthesis of N (4-methoxyphenyl, 2-butylBenzofuran-3, 5-yl) thiophene-3-carboxamide. (DBMY-20)

Yield is 62.0% purity: 99.00%. LC-MS: m/z 434.00 [M+]. IR (KBR): 3350.1cm⁻¹(NH str), 3200.2cm⁻¹(C-H Ar str), 3050.2(C-H Ali str), 1720cm⁻¹(C=O str), ¹H NMR(DMSO-d₆): δ 1.14-1.20 (3H,t,-CH₃-); δ 1.56-2.15 (4H,m-CH₂-); δ 3.32 (2H,t,-CH₂-); δ 4.19 (3H,s,-OCH₃-); δ 4.68 (1H,s,-NH-); δ 7.22-7.78 (8H,m,-Ar-). Anal. Calcd for C₂₅H₂₃NO₄S: C, 69.26; H, 5.35; N, 3.23; O, 14.76. Found: C, 68.00; H, 5.20; N, 3.86.

Synthesis of N (4-methoxyphenyl, 2-butylBenzofuran-3, 5-yl) Pyrrole-3-carboxamide. (DBMY-21)

Yield is 68.2% purity: 98.50%. LC-MS: m/z 417.08 [M+]. IR (KBR): 3350.1cm⁻¹(NH str), 3200.2cm⁻¹(C-H Ar str), 3050.2(C-H Ali str), 1710cm⁻¹(C=O str), ¹H NMR(DMSO-d₆): δ 0.86-0.90 (3H,t,-CH₃-); δ 1.34-1.87 (4H,m-CH₂-); δ 3.15 (2H,t,-CH₂-); δ 3.79 (3H,s,-OCH₃-); δ 6.99-7.51 (10H,m,-Ar-); δ 8.03 (2H,s,-NH-); δ 8.23-8.42 (1H,s,-Ar-NH₂-). Anal. Calcd for C₂₅H₂₄N₂O₄: C, 72.10; H, 5.81; N, 6.73; O, 15.37. Found: C, 71.39; H, 6.02; N, 6.87.

Synthesis of N (4-methoxyphenyl, 2-butylBenzofuran-3, 5-yl) Furan-3-carboxamide. (DBMY-22)

Yield is 70.45% purity: 98.66%. LC-MS: m/z 418.45 [M+]. IR (KBR): 3350.1cm⁻¹(NH str), 3200.2cm⁻¹(C-H Ar str), 3050.2(C-H Ali str), 1720cm⁻¹(C=O str), ¹H NMR(DMSO-d₆): δ 0.86-0.99 (3H,t,-CH₃-); δ 1.34-1.87 (4H,m-CH₂-); δ 3.15 (2H,t,-CH₂-); δ 3.98 (3H,s,-OCH₃-); δ 6.99-7.51 (10H,m,-Ar-); δ 8.23 (1H,s,-Ar-NH-). Anal. Calcd for C₂₅H₂₃NO₅: C, 71.93; H, 5.55; N, 3.36; O, 19.16. Found: C, 72.23; H, 6.30; N, 3.80.

Synthesis of N (4-methoxyphenyl, 2-butylBenzofuran-3, 5-yl) Isoxazole-2-carboxamide. (DBMY-23)

Yield is 74.99% purity: 97.50%. LC-MS: m/z 422.42 [M+]. IR (KBR): 3350.1cm⁻¹(NH str), 3200.2cm⁻¹(C-H Ar str), 3050.2(C-H Ali str), 1710cm⁻¹(C=O str), ¹H NMR(DMSO-d₆): δ 0.86-1.00 (3H,t,-CH₃-); δ 1.34-1.87 (4H,m-CH₂-); δ 3.15 (2H,t,-CH₂-); δ 3.98 (3H,s,-OCH₃-); δ 7.22-7.81 (10H,m,-Ar-). Anal. Calcd for C₂₄H₂₄N₂O₅: C, 68.56; H, 5.75; N, 6.66; O, 19.03. Found: C, 68.24; H, 5.26; N, 6.90.










Synthesis of N (4-methoxyphenyl, 2-butylBenzofuran-3, 5-yl) Isoxazole-3-carboxamide. (DBMY-24)

Yield is 69.65% purity: 98.50%. LC-MS: m/z 418.62 [M+]. IR (KBR): 3350.1cm⁻¹(NH str), 3200.2cm⁻¹(C-H Ar str), 3050.2(C-H Ali str), 1720cm⁻¹(C=O str), ¹H NMR(DMSO-d₆): δ 0.86-1.00 (3H,t,-CH₃-); δ 1.56-1.87 (4H,m-CH₂-); δ 3.15 (2H,t,-CH₂-); δ 3.98 (3H,s,-OCH₃-); δ 6.99-7.86 (10H,m,-Ar-); δ 8.03 (1H,s,-Ar-NH-). Anal. Calcd for C₂₄H₂₂N₂O₅: C, 68.89; H, 5.30; N, 6.69; O, 19.12. Found: C, 68.60; H, 5.96; N, 7.03.

Synthesis of N (4-methoxyphenyl, 2-butylBenzofuran-3, 5-yl) Isoxazole-4-carboxamide. (DBMY-25)

Yield is 68.79% purity: 98.60%. LC-MS: m/z 419.06 [M+]. IR (KBR): 3350.1cm⁻¹(NH str), 3200.2cm⁻¹(C-H Ar str), 3050.2(C-H Ali str), 1715cm⁻¹(C=O str), ¹H NMR(DMSO-d₆): δ 0.86-1.00 (3H,t,-CH₃-); δ 1.56-1.87 (4H,m-CH₂-); δ 3.15 (2H,t,-CH₂-); δ 3.98 (3H,s,-OCH₃-); δ 7.12-7.99 (10H,m,-Ar-); 7.99 (1H,s,-Ar-NH-). Anal. Calcd for C₂₄H₂₂N₂O₅: C, 68.89; H, 5.30; N, 6.69; O, 19.12. Found: C, 68.94; H, 6.07; N, 6.30.

Table – 1 Different heterocyclic and isoxazolic carboxylic acids

Sr. No	Compound code	R
1.	DBMY20	 Thiophene-3-carboxylic acid
2.	DBMY21	 Pyrrole-3-carboxylic Acid
3.	DBMY22	 Furan-3-carboxylic acid
4.	DBMY23	 Isoxazole-2-carboxylic acid
5.	DBMY24	 Isoxazole-3-carboxylic acid
6.	DBMY25	 Isoxazole-4-carboxylic acid
7.	DBMY26	 isoxazole-5- carboxylicacid
8.	DBMY27	 3-methyl-5-isoxazolecarboxylic acid
9.	DBMY28	 4-methyl-5-isoxazolecarboxylic acid

Synthesis of N (4-methoxyphenyl, 2-butylBenzofuran-3, 5-yl) Isoxazole-5-carboxamide. (DBMY-26)

Yield is 54.21% purity: 98.90%. LC-MS: m/z 419.75 [M+]. IR (KBR): 3350.1cm⁻¹(NH str), 3200.2cm⁻¹(C-H Ar str), 3050.2(C-H Ali str), 1720cm⁻¹(C=O str)., ¹H NMR(DMSO-d₆): δ 0.86-1.00 (3H,t,-CH₃-); δ 1.56-1.87 (4H,m-CH₂-); δ 3.15 (2H,t,-CH₂-); δ 3.98 (3H,s,-OCH₃-); δ 7.12-7.89 (10H,m,-Ar-). Anal. Calcd for C₂₄H₂₂N₂O₅ C, 68.89; H, 5.30; N, 6.69; O, 19.12: Found: C, 68.00; H, 5.90; N, 6.10.

Synthesis of N(4-methoxyphenyl,2-butylBenzofuran-3,5-yl)-3-methyl-5-isoxazolecarboxamide. (DBMY-27)

Yield is 58.69% purity: 98.43%. LC-MS: m/z 434.28 [M+]. IR (KBR): 3350.1cm⁻¹(NH str), 3200.2cm⁻¹(C-H Ar str), 3050.2(C-H Ali str), ¹H NMR(DMSO-d₆): δ 0.86-1.00 (3H,t,-CH₃-); δ 1.56-1.87 (4H,m-CH₂-); δ 3.15 (2H,t,-CH₂-); δ 3.98 (3H,s,-OCH₃-); δ 7.12-7.89 (10H,m,-Ar-); 6.90 (1H,s,-Ar-NH-). Anal. Calcd for C₂₄H₂₂N₂O₅ C, 69.43; H, 5.59; N, 6.48; O, 18.50; Found: C, 69.00; H, 6.55; N, 6.80.

Synthesis of N(4-methoxyphenyl,2-butylBenzofuran-3,5-yl)-4-methyl-5-isoxazolecarboxamide. (DBMY-28)

Yield is 68.01% purity: 98.93%. LC-MS: m/z 433.50 [M+]. IR (KBR): 3350.1cm⁻¹(NH str), 3200.2cm⁻¹(C-H Ar str), 3050.2(C-H Ali str), 1720cm⁻¹(C=O str), ¹H NMR(DMSO-d₆): δ 0.86-1.00 (3H,t,-CH₃-); δ 1.56-1.87 (4H,m-CH₂-); δ 3.15 (2H,t,-CH₂-); δ 3.98 (3H,s,-OCH₃-); δ 7.12-7.89 (8H,m,-Ar-); 8.43 (1H,s,-Ar-NH-). Anal. Calcd for C₂₄H₂₂N₂O₅ C, 69.43; H, 5.59; N, 6.48; O, 18.50; Found: C, 69.89; H, 5.87; N, 6.00.

RESULTS AND DISCUSSION**Antibacterial activity**

All the synthesized compounds were screened for their antibacterial activity against Staphylococcus aureus, Escherichia coli and pseudomonas aeruginosa. The activity was carried out using cup plate method [16-18] the zone of inhibition was measured in mm. DMF was used as a vehicle and Ciprofloxacin as standard drug for comparison. The compounds were tested at 10 mg/ml concentration and each well was loaded with 50 µg/ml among the synthesized compounds. The observations revealed that, DBMY-22, DBMY-27 and DBMY-28 with furan-3-carboxylic acid, 3-methyl-5-isoxazolecarboxylic acid and 4-methyl-5-isoxazolecarboxylic acid substitution respectively exhibited significant antibacterial activity against and it follows order S.aureus > E.coli > P.aeruginosa. The other Compounds exhibited moderate activity against the other test microorganisms. The zones of inhibition are presented in Table -2.

Table - 2 Antibacterial activity of the compounds

compound	<i>E.coli</i> (ATCC25922)		<i>S.aureus</i> (ATCC25923)		<i>P.aeruginosa</i> (ATCC27853)	
	zone of inhibition (mm)	% inhibition	zone of inhibition (mm)	% inhibition	zone of inhibition (mm)	% inhibition
DBMY-20	15	66.6	18	81.8	10	40.3
DBMY-21	12	54.5	10	40.0	15	66.6
DBMY-22	20	90.9	24	96.0	18	75.0
DBMY-23	18	81.5	16	66.6	08	33.3
DBMY-24	12	50.0	16	72.7	NS	NS
DBMY-25	10	40.3	08	33.3	12	54.5
DBMY-26	18	81.8	16	72.7	20	90.9
DBMY-27	20	90.9	18	81.8	18	75.0
DBMY-28	18	81.8	16	72.7	20	90.9
Ciprofloxacin	22	100	25	100	24	100

Antifungal activity

The synthesized compounds were evaluated invitro for antifungal activity by using standard agar disc diffusion method [16-18] against Candida albicans and Asperigillus fumigatus. DMF was used as a vehicle. The compounds were tested at 10 mg/ml concentration and each well was Loaded with 50 µg/ml of the sample. Flucanazole was used standard drug. The synthesized compounds DBMY-22, DBMY-27 and DBMY-28 with furan-3-carboxylic acid, 3-methyl-5-isoxazolecarboxylic acid and 4-methyl-5-isoxazolecarboxylic acid substitution respectively exhibited significant activity against Candida albicans and Asperigillus fumigatus respectively. The other Compounds exhibited moderate activity against the other test microorganisms. The zones of inhibition are presented in Table -3.

Table - 3 Antifungal activities of the compounds

Si No.	Compound No.	Presence/Absence of growth Candida albicans						Presence/Absence of growth Asperigillus fumigatus					
		I	II	III	IV	V	VI	I	II	III	IV	V	VI
	Dilutions												
01	DBMY21	-	-	-	+	-	-	-	-	-	+	+	+
02	DBMY22	-	-	-	+	-	-	-	-	+	-	-	-
03	DBMY23	-	-	-	-	-	-	-	-	-	+	+	+
04	DBMY24	-	+	+	+	-	-	-	-	+	-	+	+
05	DBMY25	+	+	+	+	-	+	+	+	+	+	+	+
06	DBMY26	+	+	+	+	-	+	+	-	+	+	+	+
07	DBMY27	-	-	-	-	-	-	-	-	-	-	-	-
08	DBMY28	-	-	-	-	+	-	-	-	-	-	-	+
09	FLUCANAZOLE	-	-	-	-	-	-	-	-	-	-	-	-

'+' indicate presence of growth, '-' indicates absence of growth.

Acknowledgement

Authors are thankful to principal, Mr. Paramjeet Singh Minhas, Priyadarsini College of pharmacy and research centre, for providing laboratory and Sophisticated Instrumentation Facility.

REFERENCES

- [1] M.G.Kadieva, E.T.Oganesyan, *Chemistry of Heterocyclic Compounds*, **1997**, 33(11), 1245-1258.
- [2] Rainer Schlecker, Bissersheim, Manfered Raschack, Weisercheim, *Chemical Abstracts*, **1989**, 111, col.111:57530.
- [3] K. Manna, Y. K. Agarwal, *Bioorg med chem. Lett*, **2009**, 19(10), 2688-2692.
- [4] Alejandro Urzua, Marcos C, Rezende, Carolina Masseurajano, Loretta Vasquez, *Molecules*, **2008**, 13(10), 882-891.
- [5] D. B. Aruna kumar, G. K. Prakash, M. N. Kumarasamy, B. P. Nanadheswarappa, B. S. Sheringava, K. M. Mahadevan, *Indian J Chem*, **2007**, 46B, 336-343.
- [6] Lourdes Santana, *Eur J pharm sci*, **1999**, 7,161-166.
- [7] A. Bindoli, *Pharmacol Res*, **1991**, (4), 369- 375.
- [8] Charloltz Mellin, Tullinge, *U.S. Patent Documents*, **1998**, 5,854,282.
- [9] I. Hayakawa, *Bioorg Med Chem. Lett*, **2004**, 14,455-458.
- [10] Chengzhi Zhang, Jacques Dumas, Gaetan, Ladauceur, Qian Zhao, Martin F. Hentemann, Sharad K. Verma, *U.S. Patent Documents*, **1989**, 7,420,066B2.
- [11] S. A. Galal, *Bioorg med chem. Lett*, **2009**, 19(9), 2420-2428.
- [12] Patrice C. Belangerl, John Scheiget, Dollard Des Oomeaux, Joshua Rokach, Quebec, *U.S. Patent Documents*, **1989**, 4,863,958.
- [13] M. D. Wei Sum, *American Heart Association, Inc.* **1999**, 100, 2276-2281.
- [14] Suman Malik, Suparna Ghosh, Bharti Jain, *Der Pharma Chemica*, **2010**, 2,304.
- [15] M. P. Ono M, Kung, H. F. Hou C, Kung, *Nucl Med Biol*, **2002**, 29(6), 633-642.
- [16] Yogesh D. Bommegowda et al, *J. Chem. Pharm. Res*, **2012**, 4(11), 4888-4893
- [17] S.Y. Xu, R.L. Bian, X. Chen, *Methadology of Pharmacology Experiment*, Peoples Medical Publishing house, Beijing; **1994**, 1356.
- [18] A. W. Bauer, W. M. Kirb, J. C. Shersis, M. Turck, *Am. J. Clin. Pathal*, **1996** 45(4), 493-496.
- [19] G. L. Wood, D. C. Washington, J.A. Washington, *Manual of Clinical Microbiology*, **1995**, 1327-1341.